

REVISITING BAER ELEMENTS

Amartya Goswami and Themba Dube

Received: 6 June 2024; Revised: 29 June 2024; Accepted: 3 July 2024

Communicated by Tuğçe Pekacar Çalçı

ABSTRACT. The objective of this paper is to extend certain properties observed in d -ideals of rings and d -elements of frames to Baer elements in multiplicative lattices. Additionally, we present results concerning these elements that have not been addressed in the study of d -ideals of rings. Furthermore, we introduce Baer closures and explore Baer maximal, prime, semiprime and meet-irreducible elements.

Mathematics Subject Classification (2020): 06F99, 06B23

Keywords: Multiplicative lattice, compact element, annihilator element, Baer element, Baer closure

1. Introduction

Baer-ideals were first introduced in [28] for commutative Baer rings, where they were defined as the kernels of Baer ring homomorphisms. Subsequently, in [19], the study of Baer ideals was extended to semiprime commutative rings. In [10], the concept of such ideals was generalized to any commutative ring and termed “ B -ideals”. The term “ d -ideal” in rings, not necessarily commutative, was introduced in [22], with the terminology adapted from the series of papers [17,18,25] focusing on d -ideals of Riesz spaces. Furthermore, in [5], a study was conducted on some properties of d -ideals in rings $C(X)$, referred to as z^0 -ideals. Also, d -ideals have been examined for reduced rings under the same name in [6].

The initial effort to extend d -ideals to “ d -elements” in lattices was undertaken in [21] for frames, once again borrowing terminology from [17]. Subsequent research on d -elements can be found in [15] and [9], where the second focused on maximal d -elements. It was in [4] where d -elements were further extended to encompass more general types of multiplicative lattices, not necessarily frames and labelled Baer elements.

Since the introduction of Baer elements in [4] for multiplicative lattices, several studies have been conducted on d -ideals in rings and d -elements in frames, as can be seen above. All of these studies may warrant scrutiny for further extensions to

multiplicative lattices in the sense of [4]. Furthermore, there are several properties of d -ideals of rings studied in earlier literature that have not been fully captured in their extended form in [4]. Our modest aim in this paper is to extend some of those results to Baer elements. Moreover, we shall include some aspects of Baer elements that, to the best of our knowledge, have not even been studied for d -ideals in rings per se.

Let us now briefly describe the content of this paper. In Section 2, we gather the multiplicative lattice theoretic machinery needed throughout this paper. All of the material presented is standard, with references including [2,3,11,12,29,30].

In Section 3, we present various equivalent characterizations and basic properties of Baer elements. We provide a necessary and sufficient condition for the closedness of Baer elements under finite products (Theorem 3.7), characterize lattices where every Baer element is a z -element (Theorem 3.14), characterize domains (Theorem 3.17) and discuss several relations between prime, maximal and Baer elements (Proposition 3.18). We also give a sufficient condition for the contraction of Baer elements under a multiplicative lattice homomorphism (Theorem 3.12).

In Section 4, we introduce Baer closures and discuss some properties of them (Proposition 4.2). We provide some equivalent criteria on closedness of Baer elements under joins (Proposition 4.4). We view Baer closures as nuclei of various types and apply them in studying some distinguished classes of Baer elements, namely maximal, prime, semiprime and meet-irreducible.

2. Preliminaries

From [12], recall that a *multiplicative lattice* is a complete lattice $(L, \leq, 0, 1)$ endowed with an associative, commutative multiplication (denoted by \cdot), which distributes over arbitrary joins and has 1 as multiplicative identity. Note that in the sense of [23], a multiplicative lattice is a commutative, unital quantale. For brevity, we shall write xy for $x \cdot y$ and x^n for $x \cdot \dots \cdot x$ (repeated n times). We say x is *strictly below* y , denoted by $x < y$, whenever $x \leq y$ and $x \neq y$. Recall that a map $\phi: L \rightarrow L'$ is said to be a *multiplicative lattice homomorphism* if ϕ preserves \leq , binary joins, binary meets and multiplication.

For our later purposes, in the following lemma, we collect some elementary properties of multiplication of multiplicative lattices.

Lemma 2.1. *In a multiplicative lattice L , the following hold.*

- (1) $xy \leq x$ for all $x, y \in L$.

- (2) $xy \leq x \wedge y$ for all $x, y \in L$.
- (3) $x0 = 0$ for all $x \in L$.
- (4) If $x \leq y$, then $xz \leq yz$ for all $x, y, z \in L$.
- (5) If $x \leq y$ and $u \leq v$, then $xu \leq yv$ for all $x, y, u, v \in L$.

Let us now recall a few definitions. Given an element a in L , an element b in L is called *minimal above a* if $a \leq b$ and for any element $c \in L$ with the property $a \leq c \leq b$ implies that $c = b$. An element a in L is called *nilpotent* if we have $a^n = 0$ for some $n \in \mathbb{N}^+$ (where \mathbb{N}^+ is the set of positive natural numbers) and L is said to be *reduced* if 0 is the only nilpotent element in L . An element c in L is called *compact*, if whenever $\{x_\lambda\}_{\lambda \in \Lambda} \subseteq L$ and $c \leq \bigvee_{\lambda \in \Lambda} x_\lambda$, we have $c \leq \bigvee_{i=1}^n x_{\lambda_i}$ for some $n \in \mathbb{N}^+$ and L is said to be *compactly generated* if every element of L is the join of compact elements in L . We shall denote by $\mathfrak{C}(L)$ the set of compact elements in L .

Let us pause with the recollection of our definitions in order to declare our underlying assumptions on lattices. For the rest of the paper, we will only work with reduced, compactly generated multiplicative lattices where 1 is compact and every finite product of compact elements is compact.

An element x in L is called *proper* if $x < 1$. A proper element m in L is said to be *maximal* if, whenever $x \in L$, $m \leq x$ and $x < 1$, we have $m = x$. We shall denote by $\mathfrak{M}(L)$ the set of all maximal elements in L . The following well-known result confirms that $\mathfrak{M}(L)$ is nonempty.

Lemma 2.2. *In a multiplicative lattice L , if the top element 1 in L is compact, then for every proper element a in L , there exists an $m \in \mathfrak{M}(L)$ such that $a \leq m$.*

A proper element p in L is called *prime* if, whenever $x, y \in L$, $xy \leq p$, we have either $x \leq p$ or $y \leq p$. We shall denote by $\mathfrak{P}(L)$ the set of prime elements in L . An element p in L is called a *minimal prime* if there is no prime element q in L with $q < p$. An element q in L is said to be *semiprime* if, whenever $x \in L$ and $x^2 \leq q$, we have $x \leq q$.

Let us recall that an element s in L is called *meet-irreducible* if, whenever $x, y \in L$ and $x \wedge y \leq s$, we have either $x \leq s$ or $y \leq s$. An element x in L is said to be a *zero divisor* if there exists a non-zero element y in L such that $xy = 0$ and L is said to be a *domain* if L does not have any non-zero zero divisor.

According to [1], the *radical of an element x* in L is defined as

$$\sqrt{x} := \bigvee \{y \in L \mid y \text{ is compact, } y^n \leq x \text{ for some } n \in \mathbb{N}^+\}.$$

It is then shown that

$$\begin{aligned}\sqrt{x} &= \bigwedge \{p \in \mathfrak{P}(L) \mid p \text{ is prime over } x\} \\ &= \bigwedge \{p \in \mathfrak{P}(L) \mid p \text{ is minimal prime over } x\}\end{aligned}\tag{†}$$

and x is called a *radical element* whenever $x = \sqrt{x}$. The *Jacobson radical* of a multiplicative lattice L is defined as

$$j_L := \bigwedge \{m \mid m \in \mathfrak{M}(L)\}$$

and L is said to be *semisimple* if $j_L = 0$. Recall that for any two elements a and b in L , the *residual* of a by b is defined as

$$(a : b) := \bigvee \{x \in L \mid xb \leq a\}$$

and the *annihilator* of an element a in L is defined as

$$a^\perp := \bigvee \{x \in L \mid xa = 0\}.$$

To denote the annihilator of the annihilator a^\perp of an element a , instead of $(a^\perp)^\perp$, we shall use the notation $a^{\perp\perp}$. Let us record some well-known properties of annihilators of elements in multiplicative lattices that will be useful in sequel.

Lemma 2.3. *Suppose L is a multiplicative lattice and a, b, c are elements in L . Then the following hold.*

- (1) *If $a \leq b$, then $b^\perp \leq a^\perp$.*
- (2) *$a \leq a^{\perp\perp}$.*
- (3) *$a^{\perp\perp\perp} = a^\perp$.*
- (4) *$a^{\perp\perp} \wedge b^{\perp\perp} = (ab)^{\perp\perp}$.*
- (5) *If $a^\perp \leq b^\perp$, then $(ac)^\perp \leq (bc)^\perp$.*

We shall denote by $\mathfrak{A}(L)$ the set of all annihilators of L . That is,

$$\mathfrak{A}(L) = \{x^\perp \mid x \in L\} = \{y^{\perp\perp} \mid y \in L\}.$$

It is proper to comment that, as in frames, with the partial order inherited from L , $\mathfrak{A}(L)$ is a Boolean algebra.

3. Baer elements

According to [4, Definition 1], an element a in a multiplicative lattice L is called a *Baer element* if, whenever $c \in \mathfrak{C}(L)$ and $c \leq a$, we have $c^{\perp\perp} \leq a$. We shall denote by $\mathfrak{B}(L)$ the set of all Baer elements in L .

In the next proposition, we gather some elementary results on Baer elements.

Proposition 3.1. *In a multiplicative lattice L , then the following hold.*

- (1) *An element b in L is a Baer element if and only if whenever $u, v \in \mathfrak{C}(L)$, $u^\perp = v^\perp$ and $u \leq b$, we have $v \leq b$.*
- (2) *Every Baer element in L is a radical element.*
- (3) *If b is a Baer element in L , then so is every minimal prime element above b .*
- (4) *Every Baer element in L is the meet of all prime Baer elements above it.*
- (5) *Both 0 and 1 in L are Baer elements.*
- (6) *If $\{x_\lambda\}_{\lambda \in \Lambda}$ is a family of Baer elements in L , then so is $\bigwedge_{\lambda \in \Lambda} x_\lambda$.*

Proof. (1)–(4) See [4].

(5) For any elements $u, v \in \mathfrak{C}(L)$, let us suppose that $u^\perp = v^\perp$ and $u \leq 0$. This implies that $u = 0$ and hence $0^\perp = 1 = v^\perp$. Therefore $v = 0$. The proof of 1 is a Baer element is trivial.

(6) Assume that $\{x_\lambda\}_{\lambda \in \Lambda}$ is a family of Baer elements of L . Let us suppose $a, b \in \mathfrak{C}(L)$ with $a^\perp = b^\perp$ and $b \leq \bigwedge_{\lambda \in \Lambda} x_\lambda$. Since $\bigwedge_{\lambda \in \Lambda} x_\lambda \leq x_\lambda$ and $x_\lambda \in \mathfrak{B}(L)$ for all $\lambda \in \Lambda$, we have $a \leq x_\lambda$ for all $\lambda \in \Lambda$ and hence $a \leq \bigwedge_{\lambda \in \Lambda} x_\lambda$, implying that $\bigwedge_{\lambda \in \Lambda} x_\lambda \in \mathfrak{B}(L)$. \square

Remark 3.2. Note that in the proof of Proposition 3.1(4), the assumption that L is reduced was utilized. Therefore, in particular, this assumption will be employed whenever we apply Proposition 3.1(4).

In Proposition 3.1(1), we have seen an alternative definition of a Baer element. In the next result we shall provide another one.

Proposition 3.3. *An element x in a multiplicative lattice L is a Baer element if and only if whenever $u, v \in \mathfrak{C}(L)$, $u^\perp \leq v^\perp$ and $u \leq x$, we have $v \leq x$.*

Proof. Let x be a Baer element in L . Suppose that $u, v \in \mathfrak{C}(L)$, $u^\perp \leq v^\perp$ and $u \leq x$. Since x is a Baer element and u compact, $u^{\perp\perp} \leq x$. Using this, (1) and (2) from Lemma 2.3, we obtain

$$v \leq v^{\perp\perp} \leq u^{\perp\perp} \leq x.$$

Thanks to Proposition 3.1(1), the converse is obvious. \square

In the context of frames, there is another well-known definition of a d -element (see [21, Definition and Remarks 5.1.]), which can also be extended to a Baer element in a multiplicative lattice. Furthermore, this definition coincides to one of the (equivalent) definitions mentioned earlier. We summarize all these facts in the following proposition.

Proposition 3.4. *The following are equivalent for an element d in a multiplicative lattice L .*

- (1) d is a Baer element.
- (2) $d = \bigvee \{c^{\perp\perp} \mid c \leq d, c \in \mathfrak{C}(L)\}$.

Proof. (1) \Rightarrow (2): Since L is compactly generated, $d = \bigvee \{c \in \mathfrak{C}(L) \mid c \leq d\}$. Now if d is a Baer element, then $v^{\perp\perp} \leq d$, whenever $v \in \mathfrak{C}(L)$ and $v \leq d$. Thus, since $w \leq w^{\perp\perp}$ for all $w \in L$,

$$d = \bigvee \{c \in \mathfrak{C}(L) \mid c \leq d\} \leq \bigvee \{c^{\perp\perp} \mid c \leq d, c \in \mathfrak{C}(L)\} \leq d,$$

which proves that $d = \bigvee \{c^{\perp\perp} \mid c \leq d, c \in \mathfrak{C}(L)\}$.

(2) \Rightarrow (1): Let us suppose that $d = \bigvee \{c^{\perp\perp} \mid c \leq d, c \in \mathfrak{C}(L)\}$. Then, whenever $v \in \mathfrak{C}(L)$ and $v \leq d$, we have $v^{\perp\perp} \leq d$, which implies d is a Baer element. \square

Remark 3.5. Let us collect all the characterizations of Baer elements that we have so far. In a multiplicative lattice L , the following are equivalent for an element $b \in L$:

- $b \in \mathfrak{B}(L)$.
- Whenever $u, v \in \mathfrak{C}(L)$, $u^\perp = v^\perp$ and $u \leq b$, we have $v \leq b$.
- Whenever $u, v \in \mathfrak{C}(L)$, $u^\perp \leq v^\perp$ and $u \leq b$, we have $v \leq b$.
- $b = \bigvee \{c^{\perp\perp} \mid c \leq b, c \in \mathfrak{C}(L)\}$.

Later on, in Proposition 4.2(2), we shall see one more (equivalent) characterization of a Baer element.

In Proposition 3.1(6), we have shown that Baer elements are closed under arbitrary meets. The next result will give us a partial converse of that and extend [26, Proposition 2.14].

Proposition 3.6. *Let $\{a_i\}_{i=1}^n$ be a set of elements in a multiplicative lattice L with the property that $a_i \vee a_j = 1$ whenever $i \neq j$. If $\bigwedge_{i=1}^n a_i$ is a Baer element in L , then each a_i ($1 \leq i \leq n$) is also a Baer element in L .*

Proof. Suppose u and v are compact elements in L , $u^\perp \leq v^\perp$ and $u \leq a_j$ for some $j \in \{1, \dots, n\}$. It is easy to see that $a_j \vee \bigwedge_{k=1, k \neq j}^n a_k = 1$, from which we obtain

$$va_j \vee \bigwedge_{k=1, k \neq j}^n va_k = v.$$

Observe that $u^\perp \leq v^\perp$ implies $(ua_k)^\perp \leq (va_k)^\perp$, for any $k \neq j$. Since $ua_k \leq \bigwedge_{k=1}^n a_k$ and $\bigwedge_{k=1}^n a_k \in \mathfrak{B}(L)$,

$$\bigwedge_{k=1, k \neq j}^n va_k \leq \bigwedge_{i=1}^n a_k,$$

and hence $\bigwedge_{k=1, k \neq j}^n va_k \leq a_j$. Obviously $va_j \leq a_j$. Therefore $v \leq a_j$, proving that $a_j \in \mathfrak{B}(L)$. \square

We shall now inquire when Baer elements are closed under finite products and the following theorem provides us with a necessary and sufficient condition for that.

Theorem 3.7. *The product of two Baer elements in a multiplicative lattice L is a Baer element if and only if $c^{\perp\perp}c^{\perp\perp} = c^{\perp\perp}$ for all compact elements c in L .*

Proof. If L has no non-zero Baer elements, then $L = \{0\}$ (cf. Proposition 3.1(5)). Let us suppose that a and b are two Baer elements in L . Suppose $c \leq ab$ for some compact element c in L . This implies that $c \leq a$ and $c \leq b$. Since a and b are Baer elements, we have $c^{\perp\perp} \leq a$ and $c^{\perp\perp} \leq b$. Applying the hypothesis and Lemma 2.1, we obtain

$$c^{\perp\perp} = c^{\perp\perp}c^{\perp\perp} \leq ab,$$

implying ab is a Baer element. Conversely, let us suppose that c be a compact element in L . Since $c \leq c^{\perp\perp}$, we have $c^2 \leq (c^{\perp\perp})^2$. Moreover, $(c^{\perp\perp})^2$ is a Baer element by our assumption. Using all these and Lemma 2.3(4), we have the desired identity from the following:

$$c^{\perp\perp} = c^{\perp\perp} \wedge c^{\perp\perp} = (cc)^{\perp\perp} \leq (c^{\perp\perp})^2 = c^{\perp\perp}c^{\perp\perp} \leq c^{\perp\perp}. \quad \square$$

Definition 3.8. We shall say a multiplicative lattice L is a *B-multiplicative lattice* if $(c^{\perp\perp})^2 = c^{\perp\perp}$ holds for all compact elements c in L .

Our next goal is to obtain some examples of Baer elements using residuals.

Proposition 3.9. *In a multiplicative lattice L , the following hold.*

- (1) *If a is a Baer element in L , then so is $(a : b)$ for any element b in L .*
- (2) *For every element a in L , the element a^\perp is a Baer element.*

- (3) If $b, \{b_i\}_{i \in I}, c$ are elements in L and $a, \{a_j\}_{j \in J}$ are Baer elements in L , then $((a : b) : c), (a : bc), ((a : c) : b), (\bigwedge_{j \in J} a_j : b), \bigwedge_{j \in J} (a_j : b), (a : \bigvee_{i \in I} b_i)$, and $\bigwedge_{i \in I} (a : b_i)$ are Baer elements in L .

Proof. (1) Let us suppose u and v are compact elements in L , $u^\perp \leq v^\perp$ and $u \leq (a : b)$. This implies that

$$bu \leq (a : b)b \leq a.$$

Since $(bu)^\perp \leq (bv)^\perp$ by Lemma 2.3(5) and $a \in \mathfrak{B}(L)$, we have $bv \leq a$, which implies that $v \leq (a : b)$.

(2) Although the proof follows from (1) and Proposition 3.1(5), however, we give another proof. Suppose $v \in \mathfrak{C}(L)$ and $v \leq a^\perp$. Applying (1) and (3) from Lemma 2.3, we have $v^{\perp\perp} \leq a^\perp$. Hence $a^\perp \in \mathfrak{B}(L)$.

(3) Follows from (1). □

We next wish to show that certain types of multiplicative lattice homomorphisms “contract” Baer elements to Baer elements. Since we are going to impose certain conditions on the homomorphisms, we first show by a simple example that the conditions we will impose are not so stringent as to make them isomorphisms.

Example 3.10. Let $\mathbf{2}$ be the two-element chain $\{0, 1\}$ and $\mathbf{3} = \{0, m, 1\}$ be the three-element chain; both viewed as multiplicative lattices with product given by \wedge . Let $\phi: \mathbf{2} \rightarrow \mathbf{3}$ be the unique homomorphism. Then ϕ has the following properties:

- ϕ maps compact elements to compact elements.
- ϕ is injective, but not an isomorphism.
- For any compact c in the domain of ϕ , $\phi(c^\perp) = \phi(c)^\perp$.

Definition 3.11. We call a multiplicative lattice homomorphism $\phi: L \rightarrow M$ *strong* if it is injective, it maps compact elements to compact elements and it commutes with annihilation of compact elements in the sense that for any compact $c \in L$, we have $\phi(c^\perp) = \phi(c)^\perp$.

Theorem 3.12. *If $\phi: L \rightarrow M$ is a strong homomorphism, then $\phi^{-1}[\mathfrak{B}(M)] \subseteq \mathfrak{B}(L)$.*

Proof. Let $x \in \phi^{-1}[\mathfrak{B}(M)]$ and consider any compact elements c and d in L with $c^\perp = d^\perp$ and $c \leq x$. We aim to show that $d \leq x$, which, by Proposition 3.1(1), will prove that x is a Baer element. Since ϕ is strong, $\phi(c)$ and $\phi(d)$ are compact element of M . Moreover, $\phi(c) \leq \phi(x)$. Since $x \in \phi^{-1}[\mathfrak{B}(M)]$, we have $\phi(x) \in \mathfrak{B}(M)$. It

follows that $\phi(d) \leq \phi(x)$, whence $d \leq x$ since ϕ is injective. Thus, $x \in \mathfrak{B}(L)$, as desired. \square

In terms of similarity, one of the closest classes of elements to Baer elements is that of z -elements. To define a z -element, we need the confirmation of existence of maximal elements in multiplicative lattices, which follows from Lemma 2.2. For an a in L , let us define

$$\mathcal{M}_a := \{m \in \mathfrak{M}(L) \mid a \leq m\} \quad \text{and} \quad \mathfrak{m}_a := \bigwedge \mathcal{M}_a.$$

According to [20], an element x in L is called a z -element if, whenever $a, b \in L$, $\mathcal{M}_a = \mathcal{M}_b$ and $b \leq x$, we have $a \leq x$.

Remark 3.13. It can be shown (see [20, Lemma 2.10]) that the following three statements are equivalent in a multiplicative lattice L (cf. Remark 3.5):

- An element x is a z -element in L .
- Whenever $a, b \in L$, $\mathcal{M}_a \supseteq \mathcal{M}_b$ and $b \leq x$, we have $a \leq x$.
- Whenever $a, b \in L$, $\mathfrak{m}_b \leq \mathfrak{m}_a$ and $a \leq x$, we have $b \leq x$.

In the next theorem, we shall characterize those multiplicative lattices in which every Baer element is a z -element. This result is well-known in rings (*e.g.*, see [26, Proposition 2.9]).

Theorem 3.14. *In a multiplicative lattice L , every Baer element is a z -element if and only if L is semisimple.*

Proof. Let x be a Baer element in a semisimple multiplicative lattice L . Let us suppose that $\mathfrak{m}_b \leq \mathfrak{m}_a$ and $a \leq x$. We claim that $a^\perp \leq b^\perp$. For $c \in \mathfrak{C}(L)$ with $ca = 0$ implies that $c \leq a^\perp$. Now from [20, Lemma 2.6] and the hypothesis, it follows that

$$\mathfrak{m}_{bc} \leq \mathfrak{m}_{ac} = \mathfrak{m}_0 = \mathfrak{j}_L = 0.$$

Therefore $bc = 0$ and hence $c \leq b^\perp$. Since $x \in \mathfrak{B}(L)$, we have $b \leq x$ and hence x is a z -element in L . Conversely, let us suppose that every Baer element in L is a z -element. From Proposition 3.1(5) it follows that 0 is a z -element. Now if $a \in \mathfrak{j}_L$, then we have $\mathfrak{m}_a = \mathfrak{m}_0 = 0$, which implies that $a = 0$. \square

Corollary 3.15. *In an algebraic frame, every d -element is a z -element.*

Remark 3.16. For the ring $\mathcal{R}L$ of continuous functions over a frame L , the notions of d -ideals and z -ideals are introduced in [13, Definition 4.10], and it is not hard to

see that every d -ideal in $\mathcal{R}L$ is a z -ideal. Therefore, our Theorem 3.14 also extends this result to multiplicative lattices.

We shall now obtain a characterization of domains by the absence of non-trivial Baer elements in them and the next result extends [26, Lemma 2.11].

Theorem 3.17. *A multiplicative lattice L is a domain if and only if L has no non-zero Baer element.*

Proof. Let us suppose that a is a Baer element in a domain L . This implies $a^{\perp\perp} = 0$ and since $a \leq a^{\perp\perp}$ by Lemma 2.3(2), $a = 0$. Conversely, suppose a is an element in L such that $ax = 0$ for some $x \in L$. This implies that $x \leq a^\perp$. By Proposition 3.1(2), a^\perp is a Baer element and by hypothesis $a^\perp = 0$. Therefore $x = 0$, implying that L is a domain. \square

Our next proposition is on establishing some relations between Baer, prime and maximal elements in multiplicative lattices and first three of them extend [26, Proposition 3.1].

Proposition 3.18. *In a multiplicative lattice L , the following hold.*

- (1) *If $x \in L$, $p \in \mathfrak{P}(L)$ and $x \wedge p \in \mathfrak{B}(L)$, then either $x \in \mathfrak{B}(L)$ or $p \in \mathfrak{B}(L)$.*
- (2) *If $p, q \in \mathfrak{P}(L)$ which do not belong to a chain and $p \wedge q \in \mathfrak{B}(L)$, then both $p, q \in \mathfrak{B}(L)$.*
- (3) *If $x \in L$, $m \in \mathfrak{M}(L)$ with $x \not\leq m$ and $x \wedge m \in \mathfrak{B}(L)$, then $x, m \in \mathfrak{B}(L)$.*
- (4) *If p is a prime element in L , then either p is a Baer element or the maximal Baer elements that are below p are prime Baer elements.*

Proof. (1) Let us consider the case: $x \leq p$. Then by hypothesis, $x \wedge p = x \in \mathfrak{B}(L)$ and we are done. So, let $x \not\leq p$. Then there exists an $y \in L$ with $y \leq x$ and $y \not\leq p$. Suppose u and v are two compact elements in L with $u^\perp \leq v^\perp$ and $u \leq p$. From Lemma 2.3(5), we have $(uy)^\perp \leq (vy)^\perp$. Since $uy \leq x \wedge p$ and by hypothesis, $x \wedge p \in \mathfrak{B}(L)$, we have $vy \leq x \wedge p$ and hence $vy \leq p$. Since p is prime and $y \not\leq p$, $v \leq p$. Therefore p is a Baer element in L .

(2) Let us suppose that $x \leq q$ and $x \not\leq p$. Suppose $u, v \in \mathfrak{C}(L)$, $u^\perp \leq v^\perp$ and $u \leq p$. By Lemma 2.3(5), $(ux)^\perp \leq (vx)^\perp$. Since $ux \leq p \wedge q$ and $p \wedge q \in \mathfrak{B}(L)$, we have

$$vx \leq p \wedge q \leq p.$$

Since p is prime and $x \not\leq p$, we must have $v \leq p$ implying that p is a Baer element in L . Similarly, we can show that q is also a Baer element in L .

(3) Since $x \not\leq m$, $x \vee m = 1$ and hence by Proposition 3.6 we have the desired claim.

(4) Let us define a set

$$S := \{x \in \mathfrak{B}(L) \mid x \leq p\}.$$

Since $0 \in S$ by Proposition 3.1(5), the set S must be nonempty. Therefore, by Zorn's lemma, S has a maximal element, say m . Now, $m = p$ if and only if p is a prime Baer element. If $m < p$, then there exists a prime element q which is minimal with respect to above m and below p . By Proposition 3.1(3), q is a Baer element and hence $q \neq p$. This implies either $q = m$, hence m is a prime element, or $m < q$, contradicting maximality of m . \square

4. Baer closures and their applications

We shall now introduce a closure on a multiplicative lattice that will assign a unique Baer element to each element of the lattice and will give another alternative definition of a Baer element. Furthermore, using these closures, we shall study some distinguished classes of Baer elements in due course.

Definition 4.1. The *Baer closure* on a multiplicative lattice L is the map $\mathfrak{c}: L \rightarrow L$ defined as

$$\mathfrak{c}(a) := \bigwedge \{x \in \mathfrak{B}(L) \mid a \leq x\}.$$

In the next proposition, we shall establish some essential properties of Baer closures.

Proposition 4.2. *Suppose a and b are elements in a multiplicative lattice L . Then the following hold.*

- (1) $\mathfrak{c}(a)$ is the smallest Baer element such that $a \leq \mathfrak{c}(a)$.
- (2) $\mathfrak{c}(a) = a$ if and only if a is a Baer element.
- (3) $\mathfrak{c}(a) = 1$ if and only if $a = 1$.
- (4) $\mathfrak{c}(0) = 0$.
- (5) If $a \leq b$, then $\mathfrak{c}(a) \leq \mathfrak{c}(b)$.
- (6) $\mathfrak{c}(\mathfrak{c}(a)) = \mathfrak{c}(a)$.
- (7) $\sqrt{a} \leq \mathfrak{c}(a)$.
- (8) $\mathfrak{c}(a) = \mathfrak{c}(\sqrt{a})$.
- (9) $\mathfrak{c}(a) \vee \mathfrak{c}(b) \leq \mathfrak{c}(a \vee b) = \mathfrak{c}(\mathfrak{c}(a) \vee \mathfrak{c}(b))$.
- (10) $\mathfrak{c}(ab) = \mathfrak{c}(a \wedge b) = \mathfrak{c}(a) \wedge \mathfrak{c}(b)$.

- (11) If L is a B -multiplicative lattice, then $\mathfrak{c}(ab) = \mathfrak{c}(a)\mathfrak{c}(b)$.
(12) $\mathfrak{c}(a^n) = \mathfrak{c}(a)$, for any $n \in \mathbb{N}^+$.

Proof. (1) From Definition 4.1 it follows that $a \leq \mathfrak{c}(a)$. By Proposition 3.1(6), we know that $\mathfrak{c}(a)$ is a Baer element in L . Now suppose y is a Baer element with the property $a \leq y$. Then from Definition 4.1, $\mathfrak{c}(a) \leq y$.

(2) If a is a Baer element, then by Definition 4.1, we have $\mathfrak{c}(a) = a$. Conversely, if the identity holds, then a is a Baer element by (1).

(3) Since 1 is a Baer element by Proposition 3.1(5), the claim $\mathfrak{c}(1) = 1$ follows by (2). The converse is obvious.

(4) Follows from Proposition 3.1(5) and (2).

(5) Let us suppose that $x \in \mathfrak{B}(L)$ with $b \leq x$. Then $a \leq x$ and hence $\mathfrak{c}(a) \leq x$. Since x was an arbitrary Baer element with the property $b \leq x$, we must have $\mathfrak{c}(a) \leq \mathfrak{c}(b)$.

(6) By (1), we have $\mathfrak{c}(a) \leq \mathfrak{c}(\mathfrak{c}(a))$. Now suppose that $x \in \mathfrak{B}(L)$ with $\mathfrak{c}(a) \leq x$. Since by definition, $\mathfrak{c}(\mathfrak{c}(a))$ is the infimum of all such $x \in \mathfrak{B}(L)$, we must have $\mathfrak{c}(\mathfrak{c}(a)) \leq \mathfrak{c}(a)$.

(7) Let us suppose that $\text{Spec}_B(L)$ denotes the set of all prime Baer elements of L . Now by applying (1), (\dagger) and Proposition 3.1(4), we have

$$\begin{aligned} \sqrt{a} &= \bigwedge \{p \in \mathfrak{B}(L) \mid a \leq p\} \\ &\leq \bigwedge \{p \in \text{Spec}_B(L) \mid a \leq p\} \\ &\leq \bigwedge \{p \in \text{Spec}_B(L) \mid \mathfrak{c}(a) \leq p\} = \mathfrak{c}(a). \end{aligned}$$

(8) Applying (5), (7) and (6), we have $\mathfrak{c}(\sqrt{a}) \leq \mathfrak{c}(a)$. On the other hand, applying (5) on $a \leq \sqrt{a}$, we get $\mathfrak{c}(a) \leq \mathfrak{c}(\sqrt{a})$.

(9) Applying (1) and (5), we obtain $\mathfrak{c}(a \vee b) \leq \mathfrak{c}(\mathfrak{c}(a) \vee \mathfrak{c}(b))$, whereas applying (5), we have $\mathfrak{c}(a) \vee \mathfrak{c}(b) \leq \mathfrak{c}(a \vee b)$ and applying (6) on it gives

$$\mathfrak{c}(\mathfrak{c}(a) \vee \mathfrak{c}(b)) \leq \mathfrak{c}(\mathfrak{c}(a \vee b)) = \mathfrak{c}(a \vee b).$$

(10) Since by Lemma 2.1, we have $ab \leq a$ and $ab \leq b$, applying Lemma 2.1, Proposition 3.1(6) and (5), we obtain

$$\mathfrak{c}(ab) \leq \mathfrak{c}(a \wedge b) \leq \mathfrak{c}(a) \wedge \mathfrak{c}(b) \in \mathfrak{B}(L).$$

Therefore, by (1), to obtain the desired equalities, it suffices to show that $\mathfrak{c}(a) \wedge \mathfrak{c}(b)$ is the smallest Baer element that is above ab . Let us suppose that x is a Baer

element in L with $ab \leq x$. We shall show that

$$\mathfrak{c}(a) \wedge \mathfrak{c}(b) \leq x.$$

By $\text{Min}_L(x)$, let us denote the set of all minimal prime elements in L that are above x . By Proposition 3.1(3), if $p \in \text{Min}_L(x)$, then p is a Baer element in L . By Proposition 3.1(4), we have

$$x = \bigwedge \{p \mid p \in \text{Min}_L(x)\}.$$

Now if $p \in \text{Min}_L(x)$, then $ab \leq p$ and hence, either $a \leq p$ or $b \leq p$. Since p is a Baer element, $\mathfrak{c}(a) \wedge \mathfrak{c}(b) \leq p$. From this, $\mathfrak{c}(a) \wedge \mathfrak{c}(b) \leq x$, as required.

(11) The proof is similar to (10).

(12) Follows from (10). □

Remark 4.3. In the context of algebraic frames in which the meet of any two compact elements is compact, there is another “closure operation” (see [21, Definition and Remarks 5.1.] which is well-known for *d-elements* (that is, Baer elements in our context). To recall, for any x in a such an algebraic frame L , the *d-closure* of x is defined as the element

$$d(x) := \bigvee \{c^{\perp\perp} \mid c \leq x, c \in \mathfrak{C}(L)\}.$$

It then turns out that $d(x)$ is a *d-element* and, in fact, the smallest *d-element* above x . Since $\mathfrak{c}(x)$ is the smallest *d-element* that is above x (see Proposition 4.2(1)), we have $\mathfrak{c}(x) = d(x)$. It is worthy of note that, whereas Martínez and Zenk [21] construct the *d-closure* “from below” by taking joins, we have constructed it “from above” by taking meets. Without going into details, let us comment that this brings to mind the *socle* of a commutative ring A with identity, $\text{Soc}(A)$, since it can be built from below and also from above because

$$\begin{aligned} \text{Soc}(A) &= \sum \{J \subseteq A \mid J \text{ is a minimal ideal of } A\} \\ &= \bigcap \{E \subseteq A \mid E \text{ is an essential ideal of } A\}. \end{aligned}$$

In Section 3, we have studied closedness properties of Baer elements under meets and products. The following proposition gives some equivalent criteria on closedness of Baer elements under joins.

Proposition 4.4. *In a multiplicative lattice L , the following are equivalent.*

- (1) *If a and b are Baer elements in L , then so is $a \vee b$.*
- (2) *Whenever $a, b \in L$, we have $\mathfrak{c}(a \vee b) = \mathfrak{c}(a) \vee \mathfrak{c}(b)$.*

(3) If $\{a_\lambda\}_{\lambda \in \Lambda} \subseteq \mathfrak{B}(L)$, then $\bigvee_{\lambda \in \Lambda} a_\lambda \in \mathfrak{B}(L)$.

(4) If $\{a_\lambda\}_{\lambda \in \Lambda} \subseteq L$, then $\mathfrak{c}(\bigvee_{\lambda \in \Lambda} a_\lambda) = \bigvee_{\lambda \in \Lambda} \mathfrak{c}(a_\lambda)$.

Proof. We shall only show (1) \Rightarrow (3). The rest of the implications are straightforward. Suppose $u, v \in \mathfrak{C}(L)$, $u^\perp \leq v^\perp$ and $u \leq \bigvee_{\lambda \in \Lambda} a_\lambda$. Since L is compactly generated, there exists a finite subset $\{\lambda_1, \dots, \lambda_n\}$ of Λ such that $u \leq \bigvee_{i=1}^n a_{\lambda_i}$. Since by hypothesis $\bigvee_{i=1}^n a_{\lambda_i}$ is a Baer element, we have

$$v \leq \bigvee_{i=1}^n a_{\lambda_i} \leq \bigvee_{\lambda \in \Lambda} a_\lambda,$$

implying that $\bigvee_{\lambda \in \Lambda} a_\lambda \in \mathfrak{B}(L)$. \square

Remark 4.5. The equivalence of (1) and (3) in Proposition 4.4, extends the corresponding result on d -ideals in rings (see [14, Remark 3.2]).

We now view Baer closures as nuclei of various types. Recall from [24, p.218] that a map $\phi: L \rightarrow L$ is called a *quantic nucleus* if, whenever $a, b \in L$, we have $a \leq \phi(a)$, $\phi(\phi(a)) = \phi(a)$ and $\phi(ab) = \phi(a)\phi(b)$.

Proposition 4.6. *If L is a B -multiplicative lattice, then every Baer closure is a quantic nucleus.*

Proof. Follows from (1), (6) and (11) of Proposition 4.2. \square

According to [27, Definition 1], the map ϕ is called a *nucleus* if it satisfies $a \leq \phi(a)$, $\phi(\phi(a)) = \phi(a)$ and $\phi(a \wedge b) = \phi(a) \wedge \phi(b)$, for all a, b in L . By [8, Definition 1.1], the map ϕ is said to be a *multiplicative nucleus* if it satisfies $\phi(a) = 1$ if and only if $a = 1$ and $\phi(ab) = \phi(a \wedge b) = \phi(a) \wedge \phi(b)$ for all $a, b \in L$.

Proposition 4.7. *Every Baer closure on a multiplicative lattice is a nucleus and a multiplicative nucleus.*

Proof. The first claim follows from (1), (6) and (10) of Proposition 4.2, whereas the second follows from (3) and (10) of Proposition 4.2. \square

Here we shall gather a few results on Baer closures. The proofs of these statements are either identical or similar to those presented in [24] and [8].

Proposition 4.8. *If L is a B -multiplicative lattice, then every Baer closure as a quantic nucleus satisfies the following identities:*

$$\mathfrak{c}(ab) = \mathfrak{c}(a \mathfrak{c}(b)) = \mathfrak{c}(\mathfrak{c}(a)b) = \mathfrak{c}(\mathfrak{c}(a) \mathfrak{c}(b)),$$

for all $a, b \in L$.

Defining the join in $\mathfrak{B}(L)$ by $\bigvee'_{i \in I} a_i := \mathfrak{c}(\bigvee_{i \in I} a_i)$, we obtain the following (cf. [24, Theorem 2.1]).

Theorem 4.9. *If \mathfrak{c} is a quantic nucleus, then $\mathfrak{B}(L)$ is a multiplicative lattice via $a \odot b = \mathfrak{c}(ab)$; and $\mathfrak{c}: L \rightarrow \mathfrak{B}(L)$ is a multiplicative lattice homomorphism. Moreover, every surjective multiplicative lattice homomorphism arises (up to isomorphism) in this manner.*

Since \mathfrak{c} satisfies the property: $\mathfrak{c}(ab) = \mathfrak{c}(a) \wedge \mathfrak{c}(b)$ (see Proposition 4.2(10)), it follows that $\mathfrak{B}(L)$ is a frame with $\odot := \wedge$ and \mathfrak{c} is called a *localic nucleus* (see [24, p. 219] and [7]). Since \mathfrak{c} is a multiplicative nucleus, in fact, we get more (cf. [8, Lemma 1.2]).

Theorem 4.10. *$\mathfrak{B}(L)$ is a compact frame.*

It is well-known that to check the primeness of an element in an algebraic frame, we need only to verify the prime condition with respect to compact elements. With the applications of Baer closures, we shall now show that in multiplicative lattices, Baer elements, which need to be shown as maximal, prime, semiprime, or meet-irreducible elements, enjoy similar privileges with respect to Baer elements.

Proposition 4.11. *Suppose L is a multiplicative lattice. A Baer element m in L is maximal if and only if m is maximal among all Baer elements in L .*

Proof. The ‘only if’ part is obvious. Therefore, suppose that m is a maximal element among all Baer elements in L and let $m < a < 1$, for some $a \notin \mathfrak{B}(L)$. We need to show that m is a maximal element in L . Now we have

$$m = \mathfrak{c}(m) < a < \mathfrak{c}(a) < \mathfrak{c}(1) = 1,$$

where the first and last equalities follow respectively from Proposition 4.2(2) and Proposition 4.2(3) and the middle strict inclusion follows from Proposition 4.2(2) and the assumption that $a \notin \mathfrak{B}(L)$. Since m is a maximal element in $\mathfrak{B}(L)$ and $\mathfrak{c}(a) \in \mathfrak{B}(L)$, we have a contradiction. Thus, m is a maximal element in L . \square

The following corollary extends the second part of [9, Theorem 3.4].

Corollary 4.12. *Every maximal Baer element in a multiplicative lattice is prime.*

We shall now aim to obtain a similar result for prime Baer elements and semiprime Baer elements, but for B -multiplicative lattices. It is not known to us whether this restriction on lattices are also necessary.

Proposition 4.13. *Suppose L is a B -multiplicative lattice. Then the following hold.*

- (1) *A Baer element p in L is prime if and only if whenever $x, y \in \mathfrak{B}(L)$ and $xy \leq p$, we have either $x \leq p$ or $y \leq p$.*
- (2) *A Baer element q in L is semiprime if and only if whenever $x \in \mathfrak{B}(L)$ and $x^2 \leq q$, we have $x \leq q$.*

Proof. (1) Let us suppose that p is a Baer element satisfying the hypothesis. Now suppose $x, y \in L$ and $xy \leq p$. Then we obtain

$$\mathfrak{c}(x) \mathfrak{c}(y) \leq \mathfrak{c}(xy) \leq \mathfrak{c}(p) = p,$$

where the first inequality follows from Proposition 4.2(11), whereas the second inequality holds due to Proposition 4.2(5) and finally, the equality is due to the fact that p is a Baer element (see Proposition 4.2(2)). By hypothesis, $x \leq \mathfrak{c}(x) \leq p$ or $y \leq \mathfrak{c}(y) \leq p$. Hence p is a prime element in L . Once again, the ‘only if’ part is obvious.

(2) Similar to (1). □

We conclude the paper with a similar result as above, but for meet-irreducible elements. Thanks to Proposition 4.2(10), however, we do not need any further restriction (as in Proposition 4.13) on our lattices.

Proposition 4.14. *A Baer element s in a multiplicative lattice L is meet-irreducible if and only if whenever $x, y \in \mathfrak{B}(L)$ and $x \wedge y \leq s$, we have $x \leq s$ or $y \leq s$.*

Proof. Let us suppose that s is a Baer element in L satisfying the hypothesis. Let $b, b' \in L$ and $b \wedge b' \leq s$. Applying (10) and (5) from Proposition 4.2, we obtain

$$\mathfrak{c}(b) \wedge \mathfrak{c}(b') = \mathfrak{c}(b \wedge b') \leq \mathfrak{c}(s) = s.$$

By hypothesis, this implies that $\mathfrak{c}(b) \leq s$ or $\mathfrak{c}(b') \leq s$. Finally, by Proposition 4.2(1), we have the desired claim. The proof of the converse is trivial. □

Concluding Remarks 4.15. We conclude the paper with the following remarks regarding future research on Baer elements:

- Although Baer elements are closed under arbitrary meets, we are not aware of an explicit characterization of multiplicative lattices in which Baer elements are closed under joins.
- It would be interesting to find a necessary and sufficient condition for the contraction of Baer elements (*cf.* Theorem 3.12).

- In the statement of Proposition 4.13, we have assumed L is a B -multiplicative lattice. We do not know whether this assumption is also necessary.
- Finally, thanks to the referee, we have the following two questions:
 - It would be interesting to see how the results presented here are applied to other particular algebraic structures. For example, for a bounded distributive lattice L we can define three notions of Baer element: Baer ideal, Baer filter and Baer congruence of L . What is the relationship between these three notions?
 - Suppose that A is an algebra in a congruence modular variety and $\text{Con}(A)$ the complete lattice of its congruences. By means of commutator operation $[\cdot]$, $\text{Con}(A)$ becomes a multiplicative lattice, in a larger sense (see [16]). It is clear that one can define the notion of Baer congruence of A . What properties are preserved for them?

Acknowledgement. The authors wish to extend heartfelt gratitude to the anonymous referee for his/her thorough review and invaluable feedback, which greatly enhanced the paper's presentation.

Disclosure statement. This is no financial or non-financial interest that has arisen from the direct applications of this research. The authors report there are no competing interests to declare.

References

- [1] F. Alarcón, D. D. Anderson and C. Jayaram, *Some results on abstract commutative ideal theory*, Period. Math. Hungar., 30(1) (1995), 1-26.
- [2] D. D. Anderson, *Multiplicative Lattices*, Ph.D dissertation, University of Chicago, 1974.
- [3] D. D. Anderson, *Abstract commutative ideal theory without chain condition*, Algebra Universalis, 6(2) (1976), 131-145.
- [4] D. D. Anderson, C. Jayaram and P. A. Phiri, *Baer lattices*, Acta Sci. Math. (Szeged), 59(1-2) (1994), 61-74.
- [5] F. Azarpanah, O. A. S. Karamzadeh and A. R. Aliabad, *On z^0 -ideals in $C(X)$* , Fund. Math., 160(1) (1999), 15-25.
- [6] F. Azarpanah, O. A. S. Karamzadeh and A. R. Aliabad, *On ideals consisting entirely of zero divisors*, Comm. Algebra, 28(2) (2000), 1061-1073.
- [7] B. Banaschewski, *On certain localic nuclei*, Cah. Topol. Géom. Différ. Catég., 35(3) (1994), 227-237.

- [8] B. Banaschewski and R. Harting, *Lattice aspects of radical ideals and choice principles*, Proc. London Math. Soc., 50(3) (1985), 385-404.
- [9] P. Bhattacharjee, *Maximal d -elements of an algebraic frame*, Order, 36(2) (2019), 377-390.
- [10] M. Contessa, *A note on Baer rings*, J. Algebra, 118(1) (1988), 20-32.
- [11] R. P. Dilworth, *Abstract residuation over lattices*, Bull. Amer. Math. Soc., 44 (1938), 262-268.
- [12] R. P. Dilworth, *Abstract commutative ideal theory*, Pacific J. Math., 12 (1962), 481-498.
- [13] T. Dube, *Some ring-theoretic properties of almost P -frames*, Algebra Universalis, 60(2) (2009), 145-162.
- [14] T. Dube, *Rings in which sums of d -ideals are d -ideals*, J. Korean Math. Soc., 56(2) (2019), 539-558.
- [15] T. Dube and L. Sithole, *On the sublocale of an algebraic frame induced by the d -nucleus*, Topology Appl., 263 (2019), 90-106.
- [16] A. Facchini, C. A. Finocchiaro and G. Janelidze, *Abstractly constructed prime spectra*, Algebra Universalis, 83(1) (2022), 8 (38 pp).
- [17] C. B. Huijsmans and B. de Pagter, *On z -ideals and d -ideals in Riesz spaces I*, Nederl. Akad. Wetensch. Indag. Math., 42(2) (1980), 183-195.
- [18] C. B. Huijsmans and B. de Pagter, *On z -ideals and d -ideals in Riesz spaces II*, Nederl. Akad. Wetensch. Indag. Math., 42(4) (1980), 391-408.
- [19] C. Jayaram, *Baer ideals in commutative semiprime rings*, Indian J. Pure Appl. Math., 15(8) (1984), 855-864.
- [20] C. S. Manjarekar and A. N. Chavan, *z -elements and z_j -elements in multiplicative lattices*, Palest. J. Math., 8(1) (2019), 138-147.
- [21] J. Martinez and E. Zenk, *When an algebraic frame is regular*, Algebra Universalis, 50(2) (2003), 231-257.
- [22] G. Mason, *Prime ideals and quotient rings of reduced rings*, Math. Japon., 34(6) (1989), 941-956.
- [23] C. J. Mulvey, &, Rend. Circ. Mat. Palermo (2) Suppl., 12 (1986), 99-104.
- [24] S. B. Niefield and K. I. Rosenthal, *Constructing locales from quantales*, Math. Proc. Cambridge Philos. Soc., 104(2) (1988), 215-234.
- [25] B. de Pagter, *On z -ideals and d -ideals in Riesz spaces III*, Nederl. Akad. Wetensch. Indag. Math., 43(4) (1981), 409-422.

- [26] S. Safaeeyan and A. Taherifar, *d-ideals, fd-ideals and prime ideals*, Quaest. Math., 42(6) (2019), 717-732.
- [27] H. Simmons, *A framework for topology*, Stud. Logic Found. Math., 96 (1978), 239-251.
- [28] T. P. Speed, *A note on commutative Baer rings*, J. Austral. Math. Soc., 14 (1972), 257-263.
- [29] M. Ward, *Residuation in structures over which a multiplication is defined*, Duke Math. J., 3(4) (1937), 627-636.
- [30] M. Ward and R. P. Dilworth, *Residuated lattices*, Trans. Amer. Math. Soc., 45(3) (1939), 335-354.

Amartya Goswami (Corresponding Author)

[1] Department of Mathematics and Applied Mathematics

Faculty of Science

University of Johannesburg

Auckland Park, South Africa

[2] National Institute for Theoretical and Computational Sciences (NITheCS)

South Africa

e-mail: agoswami@uj.ac.za

Themba Dube

[1] Department of Mathematical Sciences

University of South Africa

South Africa

[2] National Institute for Theoretical and Computational Sciences (NITheCS)

South Africa

e-mail: dubeta@unisa.ac.za