

## A NOTE ON THE SOLVABILITY OF A FINITE GROUP IN WHICH EVERY NON-NILPOTENT MAXIMAL SUBGROUP IS NORMAL

Wenjing Liu, Jiangtao Shi and Yunfeng Tian

Received: 17 October 2022; Accepted: 13 December 2022

Communicated by Abdullah Harmancı

**ABSTRACT.** We provide a new and simple proof to show that a finite group in which every non-nilpotent maximal subgroup is normal is solvable.

**Mathematics Subject Classification (2020):** 20D10

**Keywords:** Non-nilpotent maximal subgroup, solvable group, normal subgroup of prime-power order

### 1. Introduction

In this paper all groups are assumed to be finite. It is known that every maximal subgroup of a group  $G$  is normal if and only if  $G$  is a nilpotent group. As a generalization, Li and Shi [1] gave a proof to show that the following result holds.

**Theorem 1.1.** [1, Theorem 1.1] *A group  $G$  with all non-nilpotent maximal subgroups being normal is solvable.*

Moreover, based on the solvability of the group  $G$  in [1, Theorem 1.1], Shi [3, Theorem 5] proved that such a group  $G$  has a Sylow tower.

In this paper, our main goal is to provide a new and simpler proof of [1, Theorem 1.1], see Section 2.

### 2. New proof of [1, Theorem 1.1]

**Proof.** We first claim that  $G$  has a normal subgroup of prime-power order.

Suppose not. We divide the following discussions into three cases.

**Case 1:** Assume that every maximal subgroup of  $G$  is nilpotent. It follows that  $G$  is either a nilpotent group or a minimal non-nilpotent group. Then one can easily get that  $G$  has a normal Sylow subgroup that has prime-power order by the structure of minimal non-nilpotent group [2, Theorem 9.1.9], a contradiction.

**Case 2:** Assume that every maximal subgroup of  $G$  is non-nilpotent. It follows that every maximal subgroup of  $G$  is normal by the hypothesis and then  $G$  is nilpotent, this contradicts that every maximal subgroup of  $G$  is non-nilpotent.

**Case 3:** Assume that  $G$  not only has nilpotent maximal subgroups but also has non-nilpotent maximal subgroups. Since  $G$  has no normal subgroup of prime-power order, every Sylow  $p$ -subgroup  $P$  of  $G$  is not normal in  $G$  for any prime divisor  $p$  of  $|G|$ , that is  $N_G(P) < G$ . Then there exists a maximal subgroup  $M$  of  $G$  such that  $N_G(P) \leq M$ . Note that every non-nilpotent maximal subgroup of  $G$  is normal and  $P$  is not normal in  $G$ , one has that  $M$  is nilpotent by the Frattini-argument. Therefore, every Sylow subgroup of  $G$  is contained in some nilpotent maximal subgroup of  $G$ .

For any nilpotent maximal subgroup  $M$  of  $G$ , if there exists a prime divisor  $q$  of  $|M|$  such that the Sylow  $q$ -subgroup  $Q_M$  of  $M$  is not a Sylow  $q$ -subgroup of  $G$ , then  $N_G(Q_M) > M$  as  $M$  being nilpotent. It follows that  $Q_M$  is a normal subgroup of  $G$  of prime-power order since  $M$  is maximal in  $G$ , a contradiction.

Next assume that every Sylow subgroup of  $M$  is also a Sylow subgroup of  $G$  for any nilpotent maximal subgroup  $M$  of  $G$ .

For the case when  $G$  has exactly one nilpotent maximal subgroup  $M$ , then  $M$  is normal in  $G$ , which implies that  $G$  has a normal Sylow subgroup, a contradiction.

For another case when  $G$  has at least two nilpotent maximal subgroups. Let  $M_1$  and  $M_2$  be any two distinct nilpotent maximal subgroups of  $G$ .

(i) Suppose  $(|M_1|, |M_2|) = 1$ . Let  $N$  be a non-nilpotent maximal subgroup of  $G$ , then  $G = M_1N = M_2N$ . One has  $|G| = \frac{|M_1||N|}{|M_1 \cap N|} = \frac{|M_2||N|}{|M_2 \cap N|}$  and then  $\frac{|M_1|}{|M_1 \cap N|} = \frac{|M_2|}{|M_2 \cap N|}$ . Note that  $(\frac{|M_1|}{|M_1 \cap N|}, \frac{|M_2|}{|M_2 \cap N|}) = 1$  by the hypothesis. It follows that  $|M_1| = |M_1 \cap N|$  and then  $M_1 \leq N$ . One has  $M_1 = N$  since  $M_1$  is maximal in  $G$ , a contradiction.

(ii) Suppose  $(|M_1|, |M_2|) > 1$  and  $|M_1| \neq |M_2|$ . Then there exists a prime  $r$  such that  $r \mid (|M_1|, |M_2|)$ . Let  $R_1$  be a Sylow  $r$ -subgroup of  $M_1$  and  $R_2$  be a Sylow  $r$ -subgroup of  $M_2$ . Since both  $R_1$  and  $R_2$  are also Sylow  $r$ -subgroups of  $G$ , there exists an  $x \in G$  such that  $R_2 = R_1^x$ . That is  $R_1^x \in \text{Syl}_r(M_2)$ . It follows that  $R_1 \in \text{Syl}_r(M_2^{x^{-1}})$ . Since  $|M_1| \neq |M_2|$ , one has  $M_1 \neq M_2^{x^{-1}}$ . Then  $N_G(R_1) \geq \langle M_1, M_2^{x^{-1}} \rangle > M_1$ . Thus  $N_G(R_1) = G$  since  $M_1$  is maximal in  $G$ , which implies that  $R_1$  is a normal Sylow subgroup of  $G$ , a contradiction.

(iii) Suppose that all nilpotent maximal subgroups of  $G$  have the same order. Let  $M$  be any nilpotent maximal subgroup of  $G$ . Since every Sylow subgroup of  $G$  is contained in some nilpotent maximal subgroup of  $G$  and all nilpotent maximal subgroups of  $G$  have the same order, one has  $|G| = |M|$ , a contradiction.

All above arguments imply that our assumption is not true. Hence  $G$  has a normal subgroup of prime-power order.

In the following let  $G_1$  be a normal subgroup of  $G$  of prime-power order. Consider the quotient group  $G/G_1$ . It is clear that every non-nilpotent maximal subgroup of  $G/G_1$  is also normal, arguing as above, one has that  $G/G_1$  has a normal subgroup  $G_2/G_1$  of prime-power order. We go on considering the quotient group  $G/G_2$ , one by one, we can obtain a normal subgroups series:  $1 = G_0 \triangleleft G_1 \triangleleft G_2 \triangleleft \cdots \triangleleft G_i \triangleleft \cdots \triangleleft G_{s-1} \triangleleft G_s = G$ , where  $s > 1$  and every quotient group  $G_i/G_{i-1}$  has prime-power order for each  $1 \leq i \leq s$ . Therefore, one has that  $G$  is solvable.  $\square$

**Acknowledgement.** The authors are very thankful to the referee who provides valuable comments for our paper.

### References

- [1] N. Li and J. Shi, *A note on a finite group with all non-nilpotent maximal subgroups being normal*, Ital. J. Pure Appl. Math., 42 (2019), 700-702.
- [2] D. J. S. Robinson, *A Course in the Theory of Groups* (Second Edition), Springer-Verlag, New York, 1996.
- [3] J. Shi, *A finite group in which all non-nilpotent maximal subgroups are normal has a Sylow tower*, Hokkaido Math. J., 48(2) (2019), 309-312.

**Wenjing Liu, Jiangtao Shi** (Corresponding Author) and **Yunfeng Tian**

School of Mathematics and Information Sciences

Yantai University

Yantai 264005, China

e-mails: lwjytu@qq.com (W. Liu)

shijt2005@163.com (J. Shi)

m18863093906@163.com (Y. Tian)