

## SIMPLE-SEPARABLE MODULES

Rachid Ech-chaouy and Rachid Tribak

Received: 19 April 2023; Revised: 20 December 2023; Accepted: 26 December 2023

Communicated by A. Çiğdem Özcan

**ABSTRACT.** A module  $M$  over a ring is called simple-separable if every simple submodule of  $M$  is contained in a finitely generated direct summand of  $M$ . While a direct sum of any family of simple-separable modules is shown to be always simple-separable, we prove that a direct summand of a simple-separable module does not inherit the property, in general. It is also shown that an injective module  $M$  over a right noetherian ring is simple-separable if and only if  $M = M_1 \oplus M_2$  such that  $M_1$  is separable and  $M_2$  has zero socle. The structure of simple-separable abelian groups is completely described.

**Mathematics Subject Classification (2020):** 16D10, 16D40, 16D80

**Keywords:** Separable module, simple-separable module, V-ring,  $\pi$ -V-ring

### 1. Introduction

Throughout this article,  $R$  is an associative ring with an identity element and all modules considered are unital right  $R$ -modules unless stated otherwise. Let  $M$  be an  $R$ -module. By  $E(M)$  we denote the injective hull of  $M$ . The notations  $N \subseteq M$  and  $N \leq M$  mean that  $N$  is a subset and  $N$  is a submodule of  $M$ , respectively. By  $\mathbb{Q}$ ,  $\mathbb{Z}$  and  $\mathbb{N}$  we denote the ring of rational numbers, the ring of integer numbers and the set of natural numbers, respectively. In 1937 [2], Baer introduced the notion of separable abelian groups to mean torsion-free abelian groups  $G$  for which every finite subset of  $G$  can be embedded in a completely decomposable direct summand of  $G$ . The first example given by Baer of groups satisfying this property was the direct product of countably many copies of  $\mathbb{Z}$ . In 1973 [9, p. 1], Fuchs called an abelian group  $G$  for which every finite subset of  $G$  can be embedded in a direct summand  $A$  of  $G$  such that  $A$  is a direct sum of groups of rank 1 a separable group. On the other hand, another version of separability was introduced in 1968 [10] by Griffith who called an abelian group  $G$  separable if every finitely generated subgroup of  $G$  is contained in a finitely generated direct summand of  $G$ . This variation of separable groups was extended by Zöschinger in 1979 [24] to the general module theoretic setting. Following Zöschinger, a module  $M$  over an arbitrary ring  $R$  is

called separable if every finitely generated submodule of  $M$  is contained in a finitely generated direct summand of  $M$ .

In this paper, we study the “simple” version of separable modules. A module  $M$  is called simple-separable if every simple submodule of  $M$  is contained in a finitely generated direct summand of  $M$ . Note that this notion can also be considered as the dual of the notion of  $\mathfrak{m}$ -coseparable modules studied in [5]. In Section 2, we present some basic properties of these modules. It is shown that the property of being simple-separable is closed under direct sums, while a direct summand of a simple-separable module may not inherit the property. We investigate the class of rings  $R$  for which every injective  $R$ -module is simple-separable. We also prove that the class of commutative rings  $R$  for which every finitely cogenerated  $R$ -module is simple-separable is precisely that of the  $\pi$ -V-rings. Moreover, we determine the structure of simple-separable abelian groups. In Section 3, we shed some light on the modules  $M$  for which every direct summand of  $M$  is simple-separable. We conclude the paper by a short section on modules  $M$  for which every proper simple submodule of  $M$  is contained in a proper finitely generated direct summand.

## 2. Simple-separable modules

**Definition 2.1.** A module  $M$  is called *simple-separable* if every simple submodule of  $M$  is contained in a finitely generated direct summand of  $M$ .

Clearly, every separable module is simple-separable. However, the converse is not true, in general. To see this, we can consider the  $\mathbb{Z}$ -module  $\mathbb{Q}$  which is simple-separable since  $\mathbb{Q}$  has no simple submodules. On the other hand,  $\mathbb{Q}$  is not separable since  $\mathbb{Q}$  has no nonzero finitely generated direct summands.

Recall that a submodule  $N$  of a module  $M$  is called *small* in  $M$  (denoted by  $N \ll M$ ) if  $M \neq N + X$  for any proper submodule  $X$  of  $M$ .

**Remark 2.2.** Let  $M$  be an  $R$ -module. It is well known that a simple submodule of  $M$  is either small in  $M$  or a direct summand of  $M$ . It follows that  $M$  is simple-separable if and only if every simple small submodule of  $M$  is contained in a finitely generated direct summand of  $M$ . For example, if  $M$  is a module with  $\text{Rad}(M) = 0$ , then  $M$  is a simple-separable module.

**Example 2.3.** (i) It is obvious that any module  $M$  with  $\text{Soc}(M) = 0$  is simple-separable.  
(ii) It is clear that finitely generated modules are simple-separable. Also, any module which is a direct sum of finitely generated submodules (e.g., a free

module) is simple-separable. On the other hand, note that a module with small radical need not be simple-separable (see Example 2.23).

- (iii) Let a module  $M = \sum_{i \in I} L_i$  such that  $\{L_i \mid i \in I\}$  is a chain of finitely generated direct summands of the module  $M$ . It is clear that  $M$  is separable and hence  $M$  is simple-separable.
- (iv) It is easily seen that for any separable module  $M_1$  and any module  $M_2$  with  $\text{Soc}(M_2) = 0$ , the module  $M = M_1 \oplus M_2$  is simple-separable.

The proof of the next result is straightforward and hence is omitted.

**Proposition 2.4.** *Let  $M$  be an indecomposable module. Then the following statements are equivalent:*

- (i)  $M$  is simple-separable;
- (ii)  $\text{Soc}(M) = 0$  or  $M$  is finitely generated.

The following corollary is an immediate consequence of Proposition 2.4.

**Corollary 2.5.** *Let  $S$  be a simple module. Then  $E(S)$  is simple-separable if and only if  $E(S)$  is a finitely generated module.*

In the following example, we present some indecomposable simple-separable modules. Moreover, we provide an example of a simple-separable module which has a factor module which is not simple-separable.

**Example 2.6.** (i) Let  $p$  be a prime number. From Proposition 2.4, it follows that the indecomposable  $\mathbb{Z}$ -module  $\mathbb{Z}(p^\infty)$  is not simple-separable but the indecomposable  $\mathbb{Z}$ -modules  $\mathbb{Q}$ ,  $\mathbb{Z}$  and  $\mathbb{Z}/p^k\mathbb{Z}$  ( $k \in \mathbb{N}$ ) are simple-separable.

(ii) Let  $p$  be a prime number. Then the  $\mathbb{Z}$ -module  $\mathbb{Z}(p^\infty)$  is not simple-separable. On the other hand, there exists a free  $\mathbb{Z}$ -module  $F$  such that  $\mathbb{Z}(p^\infty) \cong F/L$  for some submodule  $L$  of  $F$ . It is clear that  $F$  is simple-separable since  $F$  is a direct sum of cyclic submodules.

Next, we will be concerned with direct summands of simple-separable modules. We begin by providing an example which shows that being simple-separable is not preserved by taking direct summands.

**Example 2.7.** It was shown in [16, Proposition 3.3] that there is a cyclic artinian module  $M$  over a ring  $R$  and a direct summand  $N$  of  $M^{(\mathbb{N})}$  such that  $N$  has no nonzero finitely generated direct summands. Since  $M$  is artinian, its socle is essential. Therefore  $M^{(\mathbb{N})}$  has an essential socle by [1, Propositions 6.17 and 9.19]. This implies that  $\text{Soc}(N) \neq 0$ . It follows that  $N$  is not simple-separable.

A module  $M$  is called a *D3-module* if for every pair  $(M_1, M_2)$  of direct summands of  $M$  with  $M_1 + M_2 = M$ ,  $M_1 \cap M_2$  is also a direct summand of  $M$ . It is well known that quasi-projective modules are D3-modules (see [17, Proposition 4.38]).

In contrast to Example 2.7, we next exhibit some sufficient conditions under which some special direct summands of a simple-separable module inherit the property.

**Proposition 2.8.** *Let  $M$  be a simple-separable  $R$ -module such that  $M = M_1 \oplus M_2$  is a direct sum of submodules  $M_1$  and  $M_2$  where  $M_2$  is finitely generated and semisimple. Assume that one of the following conditions is satisfied:*

- (i)  $M$  is a D3-module, or
- (ii)  $M_2$  is projective.

*Then  $M_1$  is simple-separable.*

**Proof.** Note that  $M_2$  is a finite direct sum of simple submodules. Then by induction, we can assume that  $M_2$  is a simple module. Now to prove that  $M_1$  is simple-separable, assume that  $\text{Soc}(M_1) \neq 0$  and let  $S_1$  be a simple submodule of  $M_1$ . Since  $M$  is simple-separable, there exists a finitely generated direct summand  $K$  of  $M$  such that  $S_1 \subseteq K$ . If  $K \subseteq M_1$ , we are done. Suppose now that  $K$  is not contained in  $M_1$ . Then  $K + M_1 = M$  as  $M_1$  is a maximal submodule of  $M$ .

(i) Since  $M$  is a D3-module,  $K \cap M_1$  is a direct summand of  $M$ . Therefore  $K \cap M_1$  is a direct summand of  $K$ .

(ii) Note that  $M_2 \cong M/M_1 = (K + M_1)/M_1 \cong K/(K \cap M_1)$  is projective. Then  $K \cap M_1$  is a direct summand of  $K$  (see [1, Proposition 17.2]).

Hence  $K \cap M_1$  is a finitely generated direct summand of  $M_1$  such that  $S_1 \subseteq K \cap M_1$ . It follows that  $M_1$  is a simple-separable module.  $\square$

Recall that a submodule  $N$  of a module  $M$  is called *fully invariant* if  $f(N) \subseteq N$  for every endomorphism  $f$  of  $M$ . A module  $M$  is said to have the *SIP* (Summand Intersection Property) if the intersection of any two direct summands of  $M$  is again a direct summand of  $M$ .

**Proposition 2.9.** *Let  $N$  be a submodule of a simple-separable  $R$ -module  $M$ . Assume that one of the following conditions is satisfied:*

- (i)  $N$  is a direct summand of  $M$  and  $M$  has the SIP, or
- (ii)  $N$  is a direct summand of  $M$  and  $K \cap N$  is a direct summand of  $M$  for every finitely generated direct summand  $K$  of  $M$ , or
- (iii)  $N$  is a fully invariant direct summand of  $M$ , or

(iv)  $R$  is a right noetherian ring and  $N$  is a fully invariant submodule of  $M$ .

Then  $N$  is a simple-separable module.

**Proof.** Let  $N$  be a submodule of  $M$  and let  $S$  be a simple submodule of  $N$ . Since  $M$  is simple-separable, there exists a finitely generated direct summand  $K$  of  $M$  such that  $S \subseteq K$  and  $M = K \oplus K'$  for some submodule  $K'$  of  $M$ . Moreover,  $S \subseteq K \cap N$ . The proof is completed by showing that  $K \cap N$  is a direct summand of  $N$  which is finitely generated.

(i)-(ii) Suppose that  $N$  is a direct summand of  $M$ . By hypothesis,  $K \cap N$  is a direct summand of  $M$  and hence of  $K$ . Therefore  $K \cap N$  is a finitely generated direct summand of  $N$ .

To prove (iii)-(iv), note that  $N = (K \cap N) \oplus (K' \cap N)$  since  $N$  is fully invariant in  $M$ .

(iii) As  $N$  is a direct summand of  $M$ ,  $K \cap N$  is a direct summand of  $K$  and so  $K \cap N$  is finitely generated.

(iv) Since  $R$  is right noetherian,  $K$  is a noetherian module and so  $K \cap N$  is finitely generated. This proves the proposition.  $\square$

Next, we will show that being simple-separable is preserved under direct sums.

**Theorem 2.10.** *Every direct sum of simple-separable modules is simple-separable.*

**Proof.** First note that without loss of generality, we can only prove the result for a finite direct sum of simple-separable modules. Let a module  $M = M_1 \oplus M_2$  be a direct sum of simple-separable submodules  $M_1$  and  $M_2$ . Let  $S$  be a simple submodule of  $M$ . If  $S \subseteq M_i$  for some  $i \in \{1, 2\}$ , then clearly  $S$  is contained in a finitely generated direct summand of  $M$ . Now suppose that  $S \cap M_1 = S \cap M_2 = 0$ . Then  $S \oplus M_1 = M_1 \oplus [(S \oplus M_1) \cap M_2]$ . Hence  $(S \oplus M_1) \cap M_2$  is a simple submodule of  $M_2$ . Since  $M_2$  is simple-separable, there exists a finitely generated direct summand  $K_2$  of  $M_2$  such that  $(S \oplus M_1) \cap M_2 \subseteq K_2$ . Thus  $S \oplus M_1 \subseteq M_1 \oplus K_2$ . On the other hand,  $S \oplus M_2 = [(S \oplus M_2) \cap M_1] \oplus M_2$ . Hence  $(S \oplus M_2) \cap M_1$  is a simple submodule of  $M_1$ . Since  $M_1$  is simple-separable, there exists a finitely generated direct summand  $K_1$  of  $M_1$  such that  $(S \oplus M_2) \cap M_1 \subseteq K_1$ . Thus  $S \oplus M_2 \subseteq K_1 \oplus M_2$ . Therefore

$$S \subseteq (S \oplus M_1) \cap (S \oplus M_2) \subseteq (M_1 \oplus K_2) \cap (K_1 \oplus M_2) = K_1 \oplus K_2.$$

Note that  $K_1 \oplus K_2$  is a finitely generated direct summand of  $M$ . The result follows.  $\square$

**Corollary 2.11.** *Let  $N$  be a fully invariant direct summand of a module  $M$ . Then the following conditions are equivalent:*

- (i)  $M$  is simple-separable;
- (ii)  $N$  and  $M/N$  are both simple-separable modules.

**Proof.** (i)  $\Rightarrow$  (ii) First note that  $N$  is simple-separable by Proposition 2.9(iii). To prove that  $M/N$  is simple-separable, let  $U$  be a submodule of  $M$  such that  $N \subseteq U$  and  $U/N$  is simple. By hypothesis, there exists a submodule  $K$  of  $M$  such that  $M = N \oplus K$ . Then  $U = N \oplus (U \cap K)$  and  $S = U \cap K$  is simple. Since  $M$  is simple-separable, there exist submodules  $A$  and  $B$  of  $M$  such that  $M = A \oplus B$ ,  $A$  is finitely generated and  $S \subseteq A$ . As  $N$  is fully invariant in  $M$ , we have

$$M/N = [(A + N)/N] \oplus [(B + N)/N].$$

Moreover,  $U/N \subseteq (A + N)/N$  and  $(A + N)/N \cong A/(A \cap N)$  is finitely generated.

- (ii)  $\Rightarrow$  (i) This follows from Theorem 2.10.  $\square$

Recall that a module  $M$  is called *separable* if every finitely generated submodule of  $M$  is contained in a finitely generated direct summand of  $M$ . Next, we investigate simple-separable injective modules.

**Proposition 2.12.** *The following are equivalent for an injective  $R$ -module  $M$ :*

- (i)  $M$  is a simple-separable module;
- (ii) Either  $\text{Soc}(M) = 0$  or  $E(S)$  is finitely generated for any simple submodule  $S$  of  $M$ .

*If, moreover,  $R$  is right noetherian, then (i)-(ii) are equivalent to:*

- (iii)  $M = (\oplus_{i \in I} M_i) \oplus N$  such that each  $M_i$  is an indecomposable finitely generated submodule of  $M$  and  $\text{Soc}(N) = 0$ ;
- (iv)  $M = (\oplus_{i \in I} M_i) \oplus N$  such that each  $M_i$  is a finitely generated submodule of  $M$  and  $\text{Soc}(N) = 0$ ;
- (v)  $M = L \oplus N$  such that  $L$  is a separable submodule of  $M$  and  $\text{Soc}(N) = 0$ .

**Proof.** (i)  $\Rightarrow$  (ii) Assume that  $\text{Soc}(M) \neq 0$  and let  $S$  be a simple submodule of  $M$ . By (i), there exist submodules  $K$  and  $K'$  of  $M$  such that  $M = K \oplus K'$ ,  $S \subseteq K$  and  $K$  is finitely generated. Since  $K$  is injective,  $E(S)$  is a direct summand of  $K$ . Hence  $E(S)$  is finitely generated.

- (ii)  $\Rightarrow$  (i) This is immediate.

(ii)  $\Rightarrow$  (iii) Since  $M$  is injective, there exists a submodule  $N \leq M$  such that  $M = E(\text{Soc}(M)) \oplus N$ . Set  $\text{Soc}(M) = \oplus_{i \in I} S_i$  where  $S_i$  ( $i \in I$ ) are simple submodules of  $M$ . Then  $M = (\oplus_{i \in I} E(S_i)) \oplus N$  since  $R$  is right noetherian (see [1,

Proposition 18.13]). Moreover, note that each  $E(S_i)$  ( $i \in I$ ) is a finitely generated indecomposable submodule of  $M$  and  $\text{Soc}(N) = 0$ .

(iii)  $\Rightarrow$  (iv) This is obvious.

(iv)  $\Rightarrow$  (v) This follows from the fact that any module which is a direct sum of finitely generated submodules is separable.

(v)  $\Rightarrow$  (i) This follows from Theorem 2.10.  $\square$

Following Caldwell's terminology in [3], a ring  $R$  is called *hypercyclic* if each cyclic right  $R$ -module has a cyclic injective hull. It was shown in [7, Theorems 4.1 and 4.2] that any artinian principal ideal ring is hypercyclic (see also [3, Theorem 1.5]). Commutative hypercyclic rings are characterized in [3]. From Proposition 2.12, we infer that every injective module over a hypercyclic ring is simple-separable.

In the next two corollaries, we describe simple-separable injective modules over commutative domains and over right artinian rings.

**Corollary 2.13.** *Let  $R$  be a commutative domain which is not a field. Then the following are equivalent for an injective  $R$ -module  $M$ :*

- (i)  $M$  is a simple-separable  $R$ -module;
- (ii)  $\text{Soc}(M) = 0$ .

**Proof.** Let  $E$  be an injective  $R$ -module. It is clear that  $E$  is divisible and hence  $\text{Rad}(E) = E$ .

(i)  $\Rightarrow$  (ii) Suppose that  $\text{Soc}(M) \neq 0$ . Then  $M$  contains a simple submodule  $S$ . By Proposition 2.12,  $E(S)$  is finitely generated. This contradicts the fact that  $E(S)$  has no maximal submodules (see also [11, Corollary 2]).

(ii)  $\Rightarrow$  (i) This is clear.  $\square$

**Corollary 2.14.** *Let  $M$  be an injective module over a right artinian ring  $R$ . Then the following are equivalent:*

- (i)  $M$  is a simple-separable  $R$ -module;
- (ii)  $M$  is a direct sum of finitely generated submodules.

**Proof.** This follows from Proposition 2.12 and [22, Theorem 4.5].  $\square$

As exhibited in Example 2.6, for any prime number  $p$ ,  $\mathbb{Z}(p^\infty) \cong E(\mathbb{Z}/p\mathbb{Z})$  is not simple-separable. Next, we will be concerned with the class of rings  $R$  for which every injective  $R$ -module is simple-separable.

**Proposition 2.15.** *The following are equivalent for a ring  $R$ :*

- (i) Every injective  $R$ -module is simple-separable;

- (ii)  $E(S)$  is simple-separable for any simple  $R$ -module  $S$ ;
- (iii)  $E(S)$  is finitely generated for any simple  $R$ -module  $S$ .

**Proof.** (i)  $\Rightarrow$  (ii) This is immediate.

(ii)  $\Rightarrow$  (iii) By Corollary 2.5.

(iii)  $\Rightarrow$  (i) Let  $M$  be an injective  $R$ -module and let  $S$  be a simple submodule of  $M$ . By (iii),  $E(S)$  is a finitely generated direct summand of  $M$  which contains  $S$ . Therefore  $M$  is simple-separable. This completes the proof.  $\square$

Recall that a module  $M$  is said to be *finitely cogenerated* (or *finitely embedded*) if for any family of submodules  $\{N_i : i \in I\}$  in  $M$ , if  $\bigcap_{i \in I} N_i = 0$ , then  $\bigcap_{i \in J} N_i = 0$  for some finite subset  $J \subseteq I$ . This is equivalent to the fact that  $E(M) \cong E(S_1) \oplus E(S_2) \oplus \cdots \oplus E(S_k)$  for some finitely many simple modules  $S_1, S_2, \dots, S_k$  (see [14, Proposition 19.1] and [22, p. 70]).

**Corollary 2.16.** *Let  $R$  be a commutative noetherian local ring with maximal ideal  $\mathfrak{m}$ . Then the following are equivalent:*

- (i) Every injective  $R$ -module is simple-separable;
- (ii)  $E(R/\mathfrak{m})$  is finitely generated;
- (iii)  $R$  is an artinian ring.

**Proof.** (i)  $\Leftrightarrow$  (ii) This follows from Proposition 2.15.

(ii)  $\Rightarrow$  (iii) Using Proposition 2.15, we see that  $E(R/\mathfrak{m})$  is finitely generated. Since  $R$  is noetherian, it follows that every finitely cogenerated  $R$ -module is finitely generated. Thus  $R$  is an artinian ring by [23, Theorem 3].

(iii)  $\Rightarrow$  (ii) This follows by using again [23, Theorem 3].  $\square$

**Remark 2.17.** Not every two-sided artinian ring satisfies the conditions in Proposition 2.15. In fact, even a two-sided artinian ring  $R$  may have a simple right  $R$ -module  $S$  such that  $E(S)$  is not finitely generated as illustrated in an example constructed in [15, Ex. 3.34] (see also [21, Theorem 2]).

A ring  $R$  is called a left (right)  $\pi$ -V-ring if, for every simple left (right)  $R$ -module  $S$ , the injective hull  $E(S)$  is of finite length (see [12]). Note that left and right artinian PI-rings and quasi-Frobenius rings are left and right  $\pi$ -V-rings by [21, p. 372] (see also [20, Lemma 6 and Proposition 10]).

**Example 2.18.** Using Proposition 2.15, it follows that over any right  $\pi$ -V-ring  $R$  (e.g., we can take  $R$  to be any commutative artinian ring), every injective  $R$ -module is simple-separable.



Next, we characterize the class of rings  $R$  for which every finitely cogenerated  $R$ -module is simple-separable. First we need the following lemma.

**Lemma 2.19.** *Let  $S$  be a simple module. Then the following are equivalent for  $M = E(S)$ :*

- (i) *Every submodule of  $M$  is simple-separable;*
  - (ii)  *$M$  is a noetherian module.*
- If, moreover,  $R$  is commutative, then (i)-(ii) are equivalent to:*
- (iii)  *$M$  has finite length.*

**Proof.** (i)  $\Rightarrow$  (ii) Let  $U$  be a nonzero submodule of  $M$ . Note that  $M$  is a uniform module with essential socle. Then  $U$  is indecomposable and  $\text{Soc}(U) \neq 0$ . Since  $U$  is simple-separable,  $U$  is finitely generated by Proposition 2.4. Therefore  $M$  is a noetherian module.

(ii)  $\Rightarrow$  (i) This is clear.

(ii)  $\Leftrightarrow$  (iii) Clearly,  $M$  is finitely cogenerated. The equivalence follows from [23, Proposition 4].  $\square$

**Proposition 2.20.** *The following statements are equivalent for a commutative ring  $R$ :*

- (i) *Every finitely cogenerated  $R$ -module is simple-separable;*
- (ii)  *$R$  is a  $\pi$ - $V$ -ring;*
- (iii)  *$R_{\mathfrak{m}}$  is an artinian ring for every maximal ideal  $\mathfrak{m}$  of  $R$ .*

**Proof.** (i)  $\Rightarrow$  (ii) This follows by using Lemma 2.19.

(ii)  $\Rightarrow$  (iii) This follows from [21, Theorem 5].

(iii)  $\Rightarrow$  (i) Let  $M$  be a finitely cogenerated  $R$ -module. From [23, Theorem 3], we infer that  $M$  is finitely generated. Hence  $M$  is simple-separable. This completes the proof.  $\square$

The next result is presumably well known but is included for completeness.

**Lemma 2.21.** *Let  $R$  be a commutative semilocal ring such that  $R_{\mathfrak{m}}$  is an artinian ring for every maximal ideal  $\mathfrak{m}$  of  $R$ . Then  $R$  is an artinian ring.*

**Proof.** Let  $\mathfrak{m}$  be a maximal ideal of  $R$  and put  $S_{\mathfrak{m}} = R \setminus \mathfrak{m}$ . Let  $I_1 \supseteq I_2 \supseteq \dots$  be a descending chain of ideals of  $R$ . Since  $R_{\mathfrak{m}}$  is an artinian ring, there exists an integer  $n_{\mathfrak{m}} \geq 1$  such that  $S_{\mathfrak{m}}^{-1}I_{n_{\mathfrak{m}}} = S_{\mathfrak{m}}^{-1}I_{n_{\mathfrak{m}}+i}$  for each  $i \geq 1$ . But  $R$  has only finitely many maximal ideals. So let  $n$  be the maximum of all the integers  $n_{\mathfrak{m}}$ 's (where  $\mathfrak{m}$  ranges over all of the maximal ideals of  $R$ ). It follows that  $S_{\mathfrak{m}}^{-1}I_n = S_{\mathfrak{m}}^{-1}I_{n+i}$  for every

maximal ideal  $\mathfrak{m}$  of  $R$  and all  $i \geq 1$ . This implies that  $S_{\mathfrak{m}}^{-1}(I_n/I_{n+i}) = 0$  for every maximal ideal  $\mathfrak{m}$  of  $R$  and all  $i \geq 1$ . Consequently,  $I_n = I_{n+i}$  for all  $i \geq 1$ . This shows that  $R$  is artinian.  $\square$

**Corollary 2.22.** *The following are equivalent for a commutative semilocal ring  $R$ :*

- (i) *Every finitely cogenerated  $R$ -module is simple-separable;*
- (ii)  *$R$  is an artinian ring.*

**Proof.** (i)  $\Rightarrow$  (ii) This is obvious by Proposition 2.20 and Lemma 2.21.

(ii)  $\Rightarrow$  (i) This is clear by Proposition 2.20.  $\square$

In the next example we provide a module with small radical which is not simple-separable. This example also shows that both Corollaries 2.16 and 2.22 are not true, in general, if  $R$  is not a commutative ring.

**Example 2.23.** Let  $K = F(x_1, x_2, \dots)$  with  $F$  a field. Consider the field monomorphism  $\sigma : K \rightarrow K$  defined by  $\sigma(x_i) = x_{i+1}$  for all  $i$  and  $\sigma$  is equal to the identity on  $F$ . Then  $R = K \times K$  with coordinate-wise addition and multiplication  $(x, y)(x', y') = (xx', xy' + \sigma(x')y)$  is a ring with identity. It is shown in [21, p. 375] that  $R$  is a local left artinian ring with maximal left ideal  $L = \{0\} \times K$  such that the left  $R$ -module  $E(R/L)$  is not of finite length. This implies that  $E(R/L)$  is not finitely generated since  $R$  is left artinian. Now Proposition 2.12 shows that  $E(R/L)$  is not simple-separable. On the other hand, note that  $\text{Rad}(E(R/L)) \ll E(R/L)$  by [1, Corollary 15.21].

**Remark 2.24.** There exist some commutative rings which satisfy the conditions in Proposition 2.15 but do not satisfy the statements in Proposition 2.20. For example, consider the ring  $R$  constructed in [3, Example p. 42]. In fact,  $R$  is a commutative local nonartinian hypercyclic ring. So every injective  $R$ -module is simple-separable by Proposition 2.12. On the other hand, it follows from Corollary 2.22 that not every finitely cogenerated  $R$ -module is simple-separable.

The following result shows that a simple-separable module  $M$  with  $\text{Rad}(M) = M$  contains no simple submodules.

**Proposition 2.25.** *Let  $M$  be a nonzero module with  $\text{Rad}(M) = M$ . Then  $M$  is simple-separable if and only if  $\text{Soc}(M) = 0$ .*

**Proof.** ( $\Rightarrow$ ) Assume that  $\text{Soc}(M) \neq 0$  and let  $S$  be a simple submodule of  $M$ . Since  $M$  is simple-separable, there exists a finitely generated direct summand  $K$  of  $M$  such that  $S \subseteq K$  and  $M = K \oplus K'$  for some submodule  $K'$  of  $M$ . Hence  $K$

contains a maximal submodule  $U$  with  $S \subseteq U$ . It is easily seen that  $U \oplus K'$  is a maximal submodule of  $M$ , a contradiction.

( $\Leftarrow$ ) This implication is immediate.  $\square$

Let  $G$  be an abelian group. We denote the torsion subgroup of  $G$  by  $T(G)$ . For any prime number  $p$ , let  $T_p(G) = \{x \in G \mid p^n x = 0 \text{ for some non-negative integer } n\}$  which is a subgroup of  $G$  called the  $p$ -primary component of  $G$ . Note that if  $G$  is a torsion abelian group, then  $G$  is a direct sum of its  $p$ -primary components. An abelian group  $G$  is said to be a primary group (or  $p$ -group) if  $G = T_p(G)$  for some prime  $p$ .

Let  $G$  be an abelian  $p$ -group (for some prime  $p$ ),  $x \in G$ , and  $n$  be a non-negative integer. Then  $x$  is said to have *height*  $n$  if  $x$  is divisible by  $p^n$  but not by  $p^{n+1}$  (i.e.  $x \in p^n G$  but  $x \notin p^{n+1} G$ ). In this case, we write  $h(x) = n$ . If  $x$  is divisible by  $p^k$  for every non-negative integer  $k$  (i.e.  $x \in \bigcap_{k \geq 1} p^k G$ ), then  $x$  is called an *element of infinite height* and we write  $h(x) = \infty$ . If  $x$  is an element of a subgroup  $U$  of  $G$ , then we can define two heights for  $x$ . When it is necessary, we will write  $h_U(x)$  and  $h_G(x)$  for the height of  $x$  in  $U$  and  $G$ , respectively. We always have  $h_U(x) \leq h_G(x)$ .

Recall that a subgroup  $U$  of an abelian group  $G$  is called *pure* if  $nU = U \cap nG$  for every non-negative integer  $n$ . An abelian group  $G$  is said to be of *bounded* if  $nG = 0$  for some positive integer  $n$ .

In the next theorem, we determine the structure of simple-separable abelian groups. First, we give the following four lemmas. The proof of the second one is adapted from that of [13, Theorem 9] (see also [8, Corollary 27.2]).

**Lemma 2.26.** *Let  $K$  be a finitely generated subgroup of an abelian group  $G$  with  $K \subseteq T(G)$ . Then  $K$  is a direct summand of  $T(G)$  if and only if  $K$  is a direct summand of  $G$ .*

**Proof.** The sufficiency follows by modularity. Conversely, suppose that  $K$  is a direct summand of  $T(G)$ . Then  $K$  is a pure subgroup of  $T(G)$  which is itself a pure subgroup of  $G$ . Thus  $K$  is pure in  $G$ . Moreover, note that  $K$  is a direct sum of a finite number of finite cyclic abelian groups since  $K$  is finitely generated and  $K \subseteq T(G)$  (see [8, Theorem 15.5]). Hence  $K$  is bounded. Now using [13, Theorem 7], we conclude that  $K$  is a direct summand of  $G$ .  $\square$

**Lemma 2.27.** *Let  $G$  be an abelian group such that  $\bigcap_{n \geq 1} p^n T_p(G) = 0$  for every prime number  $p$ . Then every simple subgroup of  $G$  is contained in a finite cyclic primary direct summand of  $G$ .*

**Proof.** Suppose that  $\text{Soc}(G) \neq 0$  and let  $S$  be a simple subgroup of  $G$ . Then there exist a prime number  $p$  and  $0 \neq x \in G$  such that  $S = \mathbb{Z}x \cong \mathbb{Z}/p\mathbb{Z}$ . Since  $\bigcap_{n \geq 1} p^n T_p(G) = 0$ , it follows that the subgroup  $U = T_p(G)$  has no elements of infinite height. Therefore  $x$  has finite height in  $U$ . Let  $h_U(x) = m$  for some non-negative integer  $m$ . Then there exists  $y \in U$  such that  $x = p^m y$ . Put  $H = \mathbb{Z}y$ . Clearly,  $S \subseteq H$  and  $H \cong \mathbb{Z}/p^{m+1}\mathbb{Z}$  is primary. It is easily seen that the only elements of order  $p$  in  $H$  are the multiples of  $x$  by integers prime to  $p$ . So these elements have the same height in  $H$  as in  $U$ . Thus  $H$  is a pure subgroup of  $U$  by [13, Lemma 7]. Note that  $p^{m+1}H = 0$ . It follows from [13, Theorem 7] that  $H$  is a direct summand of  $U$ . But  $U$  is a direct summand of  $T(G)$ , so  $H$  is a direct summand of  $T(G)$ . From Lemma 2.26, it follows that  $H$  is a direct summand of  $G$ . This completes the proof.  $\square$

**Lemma 2.28.** *Let  $G$  be a simple-separable abelian group. Then  $\bigcap_{n \geq 1} p^n T_p(G) = 0$  for every prime number  $p$ .*

**Proof.** Assume that  $\bigcap_{n \geq 1} p^n (T_p(G)) \neq 0$  for some prime number  $p$ . Then there exists in  $\bigcap_{n \geq 1} p^n (T_p(G))$  a nonzero element  $x$  of order  $p$ . Clearly,  $\mathbb{Z}x$  is a simple subgroup of  $G$ . Since  $G$  is simple-separable, there exists a decomposition  $G = K \oplus L$  such that  $K$  is finitely generated and  $\mathbb{Z}x \subseteq K$ . Note that  $\mathbb{Z}x \subseteq T_p(K)$ . Moreover, since  $T(K)$  is finitely generated, there exists an integer  $n \geq 1$  such that  $nT(K) = 0$ . But  $T(K)$  is a pure subgroup of  $K$ , so  $T(K)$  is a direct summand of  $K$  by [13, Theorem 7]. Note that  $T_p(K)$  is a direct summand of  $T(K)$ . Then  $T_p(K)$  is a direct summand of  $K$  which is finitely generated. Therefore there exists an integer  $s \geq 1$  such that  $p^s T_p(K) = 0$ . Since  $x \in \bigcap_{n \geq 1} p^n (T_p(G))$ , we have  $x = p^s y$  for some  $y \in T_p(G)$ . Now, since  $G = K \oplus L$ ,  $y = a + b$  for some  $a \in K$  and  $b \in L$ . Clearly,  $a \in T_p(K)$ . Therefore  $p^s a \in p^s T_p(K) = 0$  and hence  $x = p^s b \in L$ . But  $x \in K$ , so  $x \in K \cap L = 0$ , a contradiction.  $\square$

**Lemma 2.29.** *Let  $G$  be a torsion abelian group. Then  $G$  is separable if and only if  $G$  is simple-separable.*

**Proof.** The necessity is obvious. Conversely, suppose that  $G$  is simple-separable and let  $A$  be a finitely generated subgroup of  $G$ . Clearly  $A = \bigoplus_{i=1}^n T_{p_i}(A)$  for some positive integer  $n$  and distinct prime numbers  $p_i$  ( $1 \leq i \leq n$ ). Note that for every  $1 \leq i \leq n$ ,  $T_{p_i}(A)$  is a finitely generated subgroup of  $T_{p_i}(G)$ . Since each  $T_{p_i}(G)$  is a fully invariant direct summand of  $G$ , it follows from Proposition 2.9 that each  $T_{p_i}(G)$  is a simple-separable abelian group. Moreover,  $\bigoplus_{i=1}^n T_{p_i}(G)$  is a direct summand of  $G$ . The proof is completed by showing that each  $T_{p_i}(A)$  is

contained in a finitely generated direct summand of  $T_{p_i}(G)$ . So there is no loss of generality in assuming that  $G$  is a  $p$ -group for some prime number  $p$ . Since  $A$  is finitely generated,  $A$  is a finite direct sum of finite cyclic subgroups. This implies that  $A$  itself is a finite group. Since  $G$  has no nonzero elements of infinite height by Lemma 2.28, it follows that the heights of the nonzero elements of  $A$  (relative to  $G$ ) are bounded. Applying [8, Corollary 27.8], we see that  $A \subseteq B$  for some bounded direct summand  $B$  of  $G$ . Note that  $B$  is a direct sum of finite cyclic subgroups by [8, Theorem 17.2]. Since  $A$  is finitely generated, there exist subgroups  $B_1$  and  $B_2$  of  $B$  such that  $B = B_1 \oplus B_2$ ,  $B_1$  is finitely generated and  $A \subseteq B_1$ . It is clear that  $B_1$  is a direct summand of  $G$ . This finishes the proof.  $\square$

The next result should be compared with [9, Proposition 65.1] which characterized reduced abelian  $p$ -groups satisfying another variation of separability.

**Theorem 2.30.** *The following are equivalent for an abelian group  $G$ :*

- (i)  $G$  is simple-separable;
- (ii) For every prime number  $p$ ,  $\bigcap_{n \geq 1} p^n(T_p(G)) = 0$  (i.e.  $T_p(G)$  has no nonzero elements of infinite height);
- (iii) Every simple subgroup of  $G$  is contained in a finite cyclic primary direct summand of  $G$ ;
- (iv)  $T(G)$  is simple-separable;
- (v)  $T(G)$  is separable.

**Proof.** (i)  $\Rightarrow$  (ii) This implication is proved in Lemma 2.28.

(ii)  $\Rightarrow$  (iii) This is clear by Lemma 2.27.

(iii)  $\Rightarrow$  (iv) This follows immediately from Lemma 2.26 and the fact that a cyclic abelian group is either torsion or torsion-free.

(iv)  $\Rightarrow$  (v) This follows from Lemma 2.29.

(v)  $\Rightarrow$  (i) This is an immediate consequence of Lemma 2.26.  $\square$

### 3. Completely simple-separable modules

Motivated by Example 2.7, we introduce the following notion.

**Definition 3.1.** A module  $M$  is called *completely simple-separable* if every direct summand of  $M$  is simple-separable.

Recall that a module  $M$  is called a *duo module* (resp., *weak duo module*) if every submodule (resp., every direct summand) of  $M$  is fully invariant (see for example [19]).

- Example 3.2.** (i) It is clear that every module  $M$  with  $\text{Soc}(M) \cap \text{Rad}(M) = 0$  is completely simple-separable.
- (ii) Every finitely generated module is completely simple-separable.
- (iii) Using Proposition 2.9(iii), we see that every simple-separable weak duo module is completely simple-separable.
- (iv) Let  $R$  be a semiperfect ring or a simple right noetherian ring or a one-sided semihereditary ring or a one-sided principal ideal ring. Then every projective  $R$ -module is a direct sum of finitely generated submodules by [18, Theorem 3] and [16, Fact 3.4, Corollary 5.5 and Proposition 6.3]. It follows that every projective  $R$ -module is completely simple-separable.

**Proposition 3.3.** *Let  $R$  be a ring and let  $M$  be a completely simple-separable  $R$ -module. Assume that  $M$  has the ascending chain condition (ACC) on finitely generated direct summands (e.g.,  $M$  is noetherian). Then  $M = M_1 \oplus M_2$  is a direct sum of submodules  $M_1$  and  $M_2$  such that  $\text{Soc}(M_1) = 0$  and  $M_2$  is finitely generated.*

**Proof.** Suppose, to the contrary, that the module  $M$  does not have such a decomposition. Then  $\text{Soc}(M) \neq 0$ . Let  $S_1$  be a simple submodule of  $M$ . Since  $M$  is simple-separable, there exists a finitely generated direct summand  $K_1$  of  $M$  such that  $S_1 \subseteq K_1$ . Let  $N_1$  be a submodule of  $M$  such that  $M = K_1 \oplus N_1$ . Note that  $N_1$  is simple-separable and  $\text{Soc}(N_1) \neq 0$ . By similar arguments as before, it follows that  $N_1 = K_2 \oplus N_2$  such that  $K_2$  is finitely generated and  $N_2$  is a simple-separable submodule with  $\text{Soc}(N_2) \neq 0$ . By continuing this process, we get a strictly ascending chain of finitely generated direct summands  $K_1 \subsetneq K_1 \oplus K_2 \subsetneq \dots$  of  $M$ . This contradicts our assumption.  $\square$

Recall that a module  $M$  is said to have finite uniform dimension if  $M$  does not contain an infinite independent set of submodules. Dually, a module  $M$  is said to have finite hollow dimension if  $M$  does not contain an infinite coindependent family of submodules; that is, for some  $n \in \mathbb{N}$ , there exists an epimorphism from  $M$  to a direct sum of  $n$  nonzero modules but no epimorphism from  $M$  to a direct sum of more than  $n$  nonzero modules (see, for example [4, p. 47]).

It is well known that a module  $M$  has ACC on direct summands if and only if  $S = \text{End}_R(M)$  has ACC on right direct summands if and only if  $S$  contains no infinite set of nonzero orthogonal idempotents (see e.g., [5, Lemma 3.12]). Next, we present some sufficient conditions for a module to satisfy ACC on direct summands.

**Remark 3.4.** Let  $R$  be a ring and let  $M$  be an  $R$ -module. Then  $M$  has the ACC on direct summands when one of the following conditions holds.

- (i)  $M$  is artinian or noetherian (see [1, Proposition 10.14]);
- (ii)  $M$  has either finite uniform dimension or finite hollow dimension (see [4, 5.3] and [14, Proposition (6.30)']);
- (iii)  $\text{End}_R(M)$  is a semilocal ring (see [4, 5.3 and Corollary 18.7]).

In the following two results, we provide more examples of completely simple-separable modules.

**Proposition 3.5.** *Every injective simple-separable  $R$ -module is completely simple-separable.*

**Proof.** This follows directly from Proposition 2.12.  $\square$

**Proposition 3.6.** *If  $G$  is a simple-separable abelian group, then so is every subgroup of  $G$ . In particular,  $G$  is completely simple-separable.*

**Proof.** Let  $G$  be a simple-separable abelian group. From Theorem 2.30, we see that  $\bigcap_{n \geq 1} p^n(T_p(G)) = 0$  for all primes  $p$ . This implies that  $\bigcap_{n \geq 1} p^n(T_p(N)) = 0$  for any subgroup  $N$  of  $G$  and for all primes  $p$ . Now the result follows by using again Theorem 2.30.  $\square$

**Proposition 3.7.** *Let  $M$  be an artinian module. Then  $M$  is completely simple-separable if and only if  $M$  is finitely generated.*

**Proof.** The sufficiency is clear. Conversely, assume that  $M$  is completely simple-separable. From Proposition 3.3, we conclude that  $M = M_1 \oplus M_2$  such that  $\text{Soc}(M_1) = 0$  and  $M_2$  is finitely generated. As  $M$  is artinian,  $\text{Soc}(M) = \text{Soc}(M_2)$  is an essential submodule of  $M$ . This yields  $M_1 = 0$ . The result follows.  $\square$

**Proposition 3.8.** *Let  $M$  be a completely simple-separable module. Then any finitely generated semisimple submodule of  $M$  is contained in a finitely generated direct summand of  $M$ .*

**Proof.** Let  $n$  be a positive integer. We will prove that every semisimple submodule of  $M$  having uniform dimension  $n$  is contained in a finitely generated direct summand of  $M$ . This is clearly true for  $n = 1$ . Now assume that  $n \geq 2$  and every semisimple submodule of  $M$  having uniform dimension  $n - 1$  is contained in a finitely generated direct summand of  $M$ . Let  $U = S_1 \oplus S_2 \oplus \cdots \oplus S_n$  be a submodule of  $M$  which is a direct sum of  $n$  simple submodules  $S_i$  ( $1 \leq i \leq n$ ). By hypothesis,  $S_1 \oplus S_2 \oplus \cdots \oplus S_{n-1}$  is contained in a finitely generated direct summand  $K$  of  $M$ . Hence  $M = K \oplus N$  for some submodule  $N$  of  $M$ . If  $S_n \subseteq K$ , then

$U \subseteq K$ . Suppose now that  $S_n \not\subseteq K$ . In this case we have  $S_n \cap K = 0$  and hence  $S_n \oplus K = K \oplus [(S_n \oplus K) \cap N]$ . Therefore  $(S_n \oplus K) \cap N$  is a simple submodule of  $N$ . Since  $N$  is simple-separable, there exists a finitely generated direct summand  $L$  of  $N$  such that  $(S_n \oplus K) \cap N \subseteq L$ . Hence  $U \subseteq K \oplus [(S_n \oplus K) \cap N] \subseteq K \oplus L$  and  $K \oplus L$  is a finitely generated direct summand of  $M$ . This completes the proof.  $\square$

The following corollary is an immediate consequence of Proposition 3.8.

**Corollary 3.9.** *If  $M$  is a completely simple-separable module such that  $\text{Soc}(M)$  is finitely generated, then  $M = N \oplus K$  is a direct sum of submodules  $N$  and  $K$  such that  $\text{Soc}(N) = 0$  and  $K$  is finitely generated.*

**Remark 3.10.** The module  $M$  of Example 2.7 shows also that an infinite direct sum of completely simple-separable modules need not be completely simple-separable.

The next result deals with a special case of direct sums of two completely simple-separable modules.

**Proposition 3.11.** *Let  $M = M_1 \oplus M_2$  be a direct sum of submodules  $M_1$  and  $M_2$  such that  $M_1$  is completely simple-separable and  $M_2$  is semisimple. Assume that one of the following conditions is satisfied:*

- (i)  $M_2$  is projective, or
- (ii)  $M_2$  is finitely generated and  $M$  is a D3-module.

*Then  $M$  is completely simple-separable.*

**Proof.** Note first that every direct summand of a D3-module is also a D3-module by [17, Lemma 4.7]. Thus by induction it is sufficient to prove (ii) when  $M_2$  is a simple module. To prove the result, let  $N$  be a direct summand of  $M$  and let  $S$  be a simple submodule of  $N$ . We need only consider two cases:

**Case 1:** Assume that  $S$  is not contained in  $M_1$ . Then  $S \oplus M_1 = M_1 \oplus [(S \oplus M_1) \cap M_2]$  is a direct summand of  $M$  since  $M_2$  is semisimple. Hence  $S$  is a direct summand of  $N$ .

**Case 2:** Assume that  $S \subseteq M_1$ . Then  $S \subseteq N \cap M_1$ . If we prove that  $N \cap M_1$  is a direct summand of  $M$ , the assertion follows. Indeed, in this case  $N \cap M_1$  is a direct summand of  $M_1$ . This implies that  $N \cap M_1$  is simple-separable since  $M_1$  is completely simple-separable. Therefore there exists a finitely generated direct summand  $K$  of  $N \cap M_1$  such that  $S \subseteq K$ . Clearly,  $K$  is a direct summand of  $N$ .



(i) Note that  $N + M_1 = M_1 \oplus [(N + M_1) \cap M_2]$  and hence  $N/(N \cap M_1) \cong (N + M_1)/M_1 \cong (N + M_1) \cap M_2$ . Since  $(N + M_1) \cap M_2$  is a direct summand of  $M_2$ ,  $(N + M_1) \cap M_2$  is projective. Therefore  $N \cap M_1$  is a direct summand of  $M$ .

(ii) Suppose that  $M_2$  is a simple module. If  $N \subseteq M_1$ , then  $N \cap M_1 = N$  is a direct summand of  $M$ . Now assume that  $N$  is not contained in  $M_1$ . Then  $N + M_1 = M$  since  $M_1$  is a maximal submodule of  $M$ . As  $M$  is a D3-module, it follows that  $N \cap M_1$  is a direct summand of  $M$ . This completes the proof.  $\square$

#### 4. Strongly simple-separable modules

In this section, we introduce the following stronger form of simple-separability.

**Definition 4.1.** A module  $M$  is called *strongly simple-separable* if every proper simple submodule of  $M$  is contained in a proper finitely generated direct summand of  $M$ .

Note that the above notion can be considered as the “simple” version of the concept of  $\mathcal{A}$ -separable modules (see [6]).

**Example 4.2.** (i) It is easily seen that for any finitely generated module  $M_1$  and any nonzero module  $M_2$  with  $\text{Soc}(M_2) = 0$ , the module  $M = M_1 \oplus M_2$  is strongly simple-separable.

(ii) Every regular module  $M$  (i.e., every cyclic submodule of  $M$  is a direct summand) is strongly simple-separable. In particular, every semisimple module is strongly simple-separable.

(iii) If  $R$  is a right V-ring, then every  $R$ -module is strongly simple-separable since every simple  $R$ -module is injective.

**Remark 4.3.** *If a module  $M$  is not finitely generated, then  $M$  is strongly simple-separable if and only if  $M$  is simple-separable.*

The proof of the following proposition is straightforward.

**Proposition 4.4.** *Let  $M$  be an indecomposable module. Then the following conditions are equivalent:*

- (i)  $M$  is strongly simple-separable;
- (ii)  $\text{Soc}(M) = 0$  or  $M$  is a simple module.

**Remark 4.5.** Let  $S$  be a simple module. From the preceding proposition, it follows that  $E(S)$  is strongly simple-separable if and only if  $S$  is an injective module.

Next, we provide an example to show that the class of simple-separable modules and the class of strongly simple-separable modules are different.

**Example 4.6.** Let  $R$  be a commutative local artinian ring which is not a field. Let  $\mathfrak{m}$  be the maximal ideal of  $R$ . Clearly,  $R$  is not a V-ring and hence the  $R$ -module  $R/\mathfrak{m}$  is not injective. Note that  $E(R/\mathfrak{m})$  is a finitely generated  $R$ -module by [23, Theorem 3]. Then  $E(R/\mathfrak{m})$  is simple-separable. On the other hand,  $E(R/\mathfrak{m})$  is not strongly simple-separable by Remark 4.5. For example, we can take the ring  $R = \mathbb{Z}/p^n\mathbb{Z}$  for some prime number  $p$  and some integer  $n \geq 2$ . Note that in this case  $S = p^{n-1}\mathbb{Z}/p^n\mathbb{Z}$  is the unique simple  $R$ -module (up to isomorphism). Moreover,  $E(S) = R$  (see [22, Theorem 6.7]).

**Proposition 4.7.** *The following are equivalent for a ring  $R$ :*

- (i) *Every  $R$ -module is strongly simple-separable;*
- (ii) *Every injective  $R$ -module is strongly simple-separable;*
- (iii) *Every finitely cogenerated  $R$ -module is strongly simple-separable;*
- (iv)  *$R$  is a right V-ring.*

**Proof.** This follows from Example 4.2(iii) and Remark 4.5.  $\square$

In the next example, we show that the strongly simple-separable property does not always transfer from a module to each of its direct summands.

**Example 4.8.** (i) Let  $M = \bigoplus_{i \geq 1} M_i$  be a direct sum of nonzero nonsimple indecomposable finitely generated submodules  $M_i$  ( $i \geq 1$ ) such that  $\text{Soc}(M_{i_0}) \neq 0$  for some  $i_0 \geq 1$  (for example, for each  $i \geq 1$ , we can take  $M_i$  to be the  $\mathbb{Z}$ -module  $\mathbb{Z}/p_i^{n_i}\mathbb{Z}$  where  $p_i$  is a prime number and  $n_i \geq 2$  is an integer). It is clear that  $M$  is strongly simple-separable. On the other hand, using Proposition 4.4, it follows that  $M_{i_0}$  is not strongly simple-separable.

(ii) We can also consider the module  $M^{(\mathbb{N})}$  given in Example 2.7. In fact, it is easily seen that  $M^{(\mathbb{N})}$  is strongly simple-separable. But  $M^{(\mathbb{N})}$  has a direct summand which is not simple-separable.

**Proposition 4.9.** *Every direct sum of strongly simple-separable modules is strongly simple-separable.*

**Proof.** The proof can be adapted from that of Theorem 2.10 by taking into account the fact that any semisimple module is strongly simple-separable.  $\square$

The following corollary is a direct consequence of Proposition 4.9.

**Corollary 4.10.** *The following conditions are equivalent for a ring  $R$ :*

- (i) *The  $R$ -module  $R_R$  is strongly simple-separable;*
- (ii) *Every free  $R$ -module is strongly simple-separable.*

In the next result, we characterize finitely generated duo strongly simple-separable modules.

**Proposition 4.11.** *Let  $M$  be a finitely generated duo  $R$ -module which is not simple. Then  $M$  is strongly simple-separable if and only if  $\text{Soc}(M) = 0$  or  $M$  is not indecomposable.*

**Proof.** To prove the necessity, assume that  $\text{Soc}(M) \neq 0$  and let  $S$  be a simple submodule of  $M$ . Since  $M$  is strongly simple-separable and  $S \neq M$ , there exists a finitely generated proper direct summand  $K$  of  $M$  such that  $S \subseteq K$ . Hence  $M$  is not indecomposable as  $K \neq 0$ . Conversely, suppose that  $M = A \oplus B$  for some proper nonzero submodules  $A$  and  $B$  of  $M$ . Let  $T$  be a simple submodule of  $M$ . Since  $M$  is duo,  $T$  is fully invariant in  $M$ . This implies that  $T = (T \cap A) \oplus (T \cap B)$ . Since  $T$  is simple, we have  $T \subseteq A$  or  $T \subseteq B$ . This proves that  $M$  is strongly simple-separable.  $\square$

Recall that a ring  $R$  is called *right duo* if the right  $R$ -module  $R_R$  is duo. The next corollaries are direct consequences of Proposition 4.11.

**Corollary 4.12.** *Let  $R$  be a right duo ring which is not a division ring. Then the  $R$ -module  $R_R$  is strongly simple-separable if and only if  $\text{Soc}(R_R) = 0$  or  $R$  has at least one non-trivial idempotent element.*

A prime ideal  $\mathfrak{p}$  of a commutative ring  $R$  is said to be an associated prime ideal of an  $R$ -module  $M$  provided  $\mathfrak{p} = \text{Ann}_R(x)$  for some nonzero element  $x$  of  $M$ . The set of associated prime ideals of  $M$  is denoted by  $\text{Ass}(M)$ .

**Corollary 4.13.** *Let  $R$  be a commutative ring which is not a field and let  $\Omega$  be the set of all maximal ideals of  $R$ . Then the  $R$ -module  $R$  is strongly simple-separable if and only if  $\text{Ass}(R) \cap \Omega = \emptyset$  or  $R$  has at least one non-trivial idempotent element.*

We finally give the structure of strongly simple-separable abelian groups.

**Proposition 4.14.** *Let  $G$  be a simple-separable abelian group. Then the following conditions are equivalent:*

- (i)  *$G$  is strongly simple-separable;*
- (ii)  *$G$  is not isomorphic to  $\mathbb{Z}/p^n\mathbb{Z}$  for every prime number  $p$  and any integer  $n \geq 2$ .*

**Proof.** (i)  $\Rightarrow$  (ii) Given a prime number  $p$  and an integer  $n \geq 2$ , it is clear that the indecomposable nonsimple  $\mathbb{Z}$ -module  $\mathbb{Z}/p^n\mathbb{Z}$  is not strongly simple-separable since  $\text{Soc}(\mathbb{Z}/p^n\mathbb{Z}) \neq 0$  (see Proposition 4.4).

(ii)  $\Rightarrow$  (i) Let  $G$  be a simple-separable abelian group which is not isomorphic to  $\mathbb{Z}/p^n\mathbb{Z}$  for every prime number  $p$  and any integer  $n \geq 2$ . If  $G$  contains no simple proper subgroups, then clearly  $G$  is strongly simple-separable. Now assume that  $G$  contains a simple proper subgroup  $S$ . Then  $S$  is isomorphic to  $\mathbb{Z}/p_0\mathbb{Z}$  for some prime number  $p_0$ . By Theorem 2.30,  $S$  is contained in a direct summand  $H$  of  $G$  with  $H \cong \mathbb{Z}/p_0^k\mathbb{Z}$  for some positive integer  $k$ . If  $k = 1$ , then  $H = S$  and hence  $H \neq G$ . Moreover, if  $k \geq 2$ , then  $H \neq G$  by (ii). This completes the proof.  $\square$

**Acknowledgement.** The authors would like to thank the referee for the valuable suggestions and comments which improved this paper.

## Declarations

**Disclosure statement:** The authors report that there are no competing interests to declare.

## References

- [1] F. W. Anderson and K. R. Fuller, *Rings and Categories of Modules*, Grad. Texts in Math., 13, Springer-Verlag, New York-Heidelberg, 1974.
- [2] R. Baer, *Abelian groups without elements of finite order*, Duke Math. J., 3(1) (1937), 68-122.
- [3] W. H. Caldwell, *Hypercyclic rings*, Pacific J. Math., 24 (1968), 29-44.
- [4] J. Clark, C. Lomp, N. Vanaja and R. Wisbauer, *Lifting Modules. Supplements and Projectivity in Module Theory*, Front. Math., Birkhäuser, Verlag, Basel, 2006.
- [5] R. Ech-chaouy, A. Idelhadj and R. Tribak, *On coseparable and  $\mathfrak{m}$ -coseparable modules*, J. Algebra Appl., 20(3) (2021), 2150031 (20 pp).
- [6] R. Ech-chaouy, A. Idelhadj and R. Tribak, *On a class of separable modules*, Asian-Eur. J. Math., 15(2) (2022), 2250034 (17 pp).
- [7] C. Faith, *On Köthe rings*, Math. Ann., 164 (1966), 207-212.
- [8] L. Fuchs, *Infinite Abelian Groups*, vol. I, Pure Appl. Math., 36, Academic Press, New York-London, 1970.
- [9] L. Fuchs, *Infinite Abelian Groups*, vol. II, Pure Appl. Math., 36, Academic Press, New York-London, 1973.

- [10] P. Griffith, *Separability of torsion free groups and a problem of J. H. C. Whitehead*, Illinois J. Math., 12 (1968), 654-659.
- [11] H. Harui, *On injective modules*, J. Math. Soc. Japan, 21 (1969), 574-583.
- [12] Y. Hirano, *On injective hulls of simple modules*, J. Algebra, 225(1) (2000), 299-308.
- [13] I. Kaplansky, *Infinite Abelian Groups*, University of Michigan Press, Ann Arbor, 1969.
- [14] T. Y. Lam, *Lectures on Modules and Rings*, Grad. Texts in Math., 189, Springer-Verlag, New York, 1999.
- [15] T. Y. Lam, *Exercises in Modules and Rings*, Probl. Books in Math., Springer, New York, 2007.
- [16] W. Wm. McGovern, G. Puninski and P. Rothmaler, *When every projective module is a direct sum of finitely generated modules*, J. Algebra, 315(1) (2007), 454-481.
- [17] S. H. Mohamed and B. J. Müller, *Continuous and Discrete Modules*, London Math. Soc. Lecture Note Ser., 147, Cambridge University Press, Cambridge, 1990.
- [18] B. J. Müller, *On semi-perfect rings*, Illinois J. Math., 14 (1970), 464-467.
- [19] A. Ç. Özcan, A. Harmanci and P. F. Smith, *Duo modules*, Glasg. Math. J., 48(3) (2006), 533-545.
- [20] M. Rayar, *On small and cosmall modules*, Acta Math. Acad. Sci. Hungar., 39(4) (1982), 389-392.
- [21] A. Rosenberg and D. Zelinsky, *Finiteness of the injective hull*, Math. Z., 70 (1958/59/1959), 372-380.
- [22] D. W. Sharpe and P. Vámos, *Injective Modules*, Cambridge Tracts in Mathematics and Mathematical Physics, Cambridge University Press, London-New York, 1972.
- [23] P. Vámos, *The dual of the notion of "finitely generated"*, J. London Math. Soc., 43 (1968), 643-646.
- [24] H. Zöschinger, *Quasi-separable und koseparable moduln über diskreten bewertungsringen*, Math. Scand., 44 (1979), 17-36.

**Rachid Ech-chaouy**

Department of Mathematics  
Faculty of Sciences  
Abdelmalek Essaâdi University  
BP. 2121 Tetouan, Morocco  
e-mail: echaourachid@yahoo.fr

**Rachid Tribak** (Corresponding Author)

Centre Régional des Métiers de l'Education et de la Formation (CRMEF-TTH)-Tanger  
Avenue My Abdelaziz, Souani  
B.P. 3117, Tangier, Morocco  
e-mail: tribak12@yahoo.com